

Some Twisted Results

M.D. Gould and T. Lekatsas

*Department of Mathematics, The University of Queensland,
Brisbane, Qld 4072, Australia
Email: tel@maths.uq.edu.au*

February 1, 2008

Abstract

The Drinfeld twist for the opposite quasi-Hopf algebra, H^{cop} is determined and is shown to be related to the (second) Drinfeld twist on a quasi-Hopf algebra. The twisted form of the Drinfeld twist is investigated. In the quasi-triangular case it is shown that the Drinfeld u operator arises from the equivalence of H^{cop} to the quasi-Hopf algebra induced by twisting H with the R -matrix. The Altschuler-Coste u operator arises in a similar way and is shown to be closely related to the Drinfeld u operator. The quasi-cocycle condition is introduced, and is shown to play a central role in the uniqueness of twisted structures on quasi-Hopf algebras. A generalisation of the dynamical quantum Yang-Baxter equation, called the quasi-dynamical quantum Yang-Baxter equation is introduced.

1 Introduction

Quasi-Hopf algebras (QHA) were introduced by Drinfeld [6] as generalisations of Hopf algebras. QHA are the underlying algebraic structures of elliptic quantum groups [8, 9, 10, 11, 14, 20] and hence have an important role in obtaining solutions to the dynamical Yang-Baxter equation. They arise in conformal field theory [3, 4], algebraic number theory [7] and in the theory of knots [1, 15, 16].

The antipode S of a Hopf algebra H is uniquely determined as the inverse of the identity map on H under the convolution product. For a quasi-Hopf algebra, the triple (S, α, β) consisting of the antipode S and canonical elements $\alpha, \beta \in H$ is termed the *quasi-antipode*. The quasi-antipode of a QHA is not unique [2, 6, 17]. However, given two QHA which differ only in their quasi-antipodes, there exists a unique invertible element $v \in H$ relating them. Moreover, to each invertible element $v \in H$ there corresponds a quasi-antipode, so that the invertible elements $v \in H$ are in bijection with the quasi-antipodes. This allows us to work with a fixed choice for the quasi-antipode (more precisely, a fixed equivalence class for the quasi-antipode). We show that the operator $v \in H$ is universal i.e. invariant under an arbitrary twist $F \in H \otimes H$. In the quasi-triangular case, the equivalence of the quasi-antipode of the opposite QHA H^{cop} and the quasi-antipode induced by twisting H with the R -matrix, gives rise to a specific form of the v operator, which we call the Drinfeld - Reshetikhin [5, 18] u operator. The u -operator introduced by Altschuler and Coste [1], arises in a similar way and is shown to be simply related to the Drinfeld - Reshetikhin u operator. In view of the invariance of the v operators these u operators are also invariant under twisting.

For a Hopf algebra H the antipode S is both an algebra and a co-algebra anti-homomorphism. In the QHA case Drinfeld has shown that the antipode S is a co-algebra anti-homomorphism only upto conjugation by a twist, F_δ (the Drinfeld twist). Assuming the antipode S is invertible with inverse S^{-1} , we show that S^{-1} is a co-algebra anti-homomorphism upto conjugation by an invertible element F_0 , which we call the second Drinfeld twist on H . The form of the Drinfeld twist for the opposite QHA H^{cop} is determined and shown to be simply related to this second Drinfeld twist. The behaviour of the Drinfeld twist F_δ under an arbitrary twist $G \in H \otimes H$ is also investigated.

The set of twists on a QHA H form a group. We study a sub-group of the group of twists on a QHA, namely those that leave the co-product $\Delta : H \rightarrow H \otimes H$ and co-associator $\Phi \in H \otimes H \otimes H$ unchanged. These twists are called *compatible twists*. Twists that leave the coassociator Φ unchanged are said to satisfy the quasi-cocycle condition. The quasi-cocycle condition is intimately related to the uniqueness of the structure obtained by twisting the quasi-bialgebra part of a QHA. In the quasi-triangular case we show that $\mathcal{R}^T \mathcal{R}$ and its powers are compatible twists.

Following on from our considerations of the quasi-cocycle condition we introduce the shifted quasi-cocycle condition on a twist $F(\lambda) \in H \otimes H$, where $\lambda \in H$ depends on one (or more) parameters. We conclude with the quasi-dynamical quantum Yang-Baxter equation (QQYBE), which is the quasi-Hopf analogue of the usual dynamical QYBE.

2 Preliminaries

We begin by recalling the definition [6] of a quasi-bialgebra.

Definition 1. A quasi-bialgebra $(H, \Delta, \epsilon, \Phi)$ is a unital associative algebra H over a field F , equipped with algebra homomorphisms $\epsilon : H \rightarrow F$ (co-unit), $\Delta : H \rightarrow H \otimes H$ (co-product) and an invertible

element $\Phi \in H \otimes H \otimes H$ (co-associator) satisfying

$$(1 \otimes \Delta)\Delta(a) = \Phi^{-1}(\Delta \otimes 1)\Delta(a)\Phi, \quad \forall a \in H, \quad (2.1)$$

$$(\Delta \otimes 1 \otimes 1)\Phi \cdot (1 \otimes 1 \otimes \Delta)\Phi = (\Phi \otimes 1) \cdot (1 \otimes \Delta \otimes 1)\Phi \cdot (1 \otimes \Phi), \quad (2.2)$$

$$(\epsilon \otimes 1)\Delta = 1 = (1 \otimes \epsilon)\Delta, \quad (2.3)$$

$$(1 \otimes \epsilon \otimes 1)\Phi = 1. \quad (2.4)$$

It follows from equations (2.2), (2.3) and (2.4) that the co-associator Φ has the additional properties

$$(\epsilon \otimes 1 \otimes 1)\Phi = 1 = (1 \otimes 1 \otimes \epsilon)\Phi.$$

We now fix the notation to be used throughout the paper. For the co-associator we follow the notation of [12, 13] and write

$$\Phi = \sum_{\nu} X_{\nu} \otimes Y_{\nu} \otimes Z_{\nu}, \quad \Phi^{-1} = \sum_{\nu} \bar{X}_{\nu} \otimes \bar{Y}_{\nu} \otimes \bar{Z}_{\nu}.$$

We adopt Sweedler's [19] notation for the co-product

$$\Delta(a) = \sum_{(a)} a_{(1)} \otimes a_{(2)}, \quad \forall a \in H$$

throughout. Since the co-product is quasi-coassociative we use the following extension of Sweedler's notation

$$\begin{aligned} (1 \otimes \Delta)\Delta(a) &= a_{(1)} \otimes \Delta(a_{(2)}) = a_{(1)} \otimes a_{(2)}^{(1)} \otimes a_{(2)}^{(2)} \\ (\Delta \otimes 1)\Delta(a) &= \Delta(a_{(1)}) \otimes a_{(2)} = a_{(1)}^{(1)} \otimes a_{(1)}^{(2)} \otimes a_{(2)}. \end{aligned} \quad (2.5)$$

In general, the summation sign is omitted from expressions, with the convention that repeated indices are to be summed over.

Definition 2. A quasi-Hopf algebra $(H, \Delta, \epsilon, \Phi, S, \alpha, \beta)$ is a quasi-bialgebra $(H, \Delta, \epsilon, \Phi)$ equipped with an algebra anti-homomorphism S (antipode) and canonical elements $\alpha, \beta \in H$ such that

$$S(X_{\nu})\alpha Y_{\nu}\beta S(Z_{\nu}) = 1 = \bar{X}_{\nu}\beta S(\bar{Y}_{\nu})\alpha \bar{Z}_{\nu}, \quad (2.6)$$

$$S(a_{(1)})\alpha a_{(2)} = \epsilon(a)\alpha, \quad a_{(1)}\beta S(a_{(2)}) = \epsilon(a)\beta, \quad \forall a \in H. \quad (2.7)$$

Throughout we assume bijectivity of the antipode S so that S^{-1} exists. The antipode equations (2.6), (2.7) imply $\epsilon(\alpha) \cdot \epsilon(\beta) = 1$ and $\epsilon(S(a)) = \epsilon(S^{-1}(a)) = \epsilon(a), \forall a \in H$. A triple (S, α, β) satisfying equations (2.6), (2.7) is called a *quasi-antipode*.

We shall need the following relations:

$$X_{\nu}a \otimes Y_{\nu}\beta S(Z_{\nu}) = a_{(1)}^{(1)}X_{\nu} \otimes a_{(1)}^{(2)}Y_{\nu}\beta S(Z_{\nu})S(a_{(2)}), \quad \forall a \in H \quad (2.8)$$

$$\begin{aligned} \Phi \otimes 1 &\stackrel{(2.2)}{=} (\Delta \otimes 1 \otimes 1)\Phi \cdot (1 \otimes 1 \otimes \Delta)\Phi \cdot (1 \otimes \Phi^{-1}) \cdot (1 \otimes \Delta \otimes 1)\Phi^{-1} \\ &= X_{\nu}^{(1)}X_{\mu}\bar{X}_{\rho} \otimes X_{\nu}^{(2)}Y_{\mu}\bar{X}_{\sigma}\bar{Y}_{\rho}^{(1)} \otimes Y_{\nu}Z_{\mu}^{(1)}\bar{Y}_{\sigma}\bar{Y}_{\rho}^{(2)} \otimes Z_{\nu}Z_{\mu}^{(2)}\bar{Z}_{\sigma}\bar{Z}_{\rho} \end{aligned} \quad (2.9)$$

$$\begin{aligned} 1 \otimes \Phi &= (1 \otimes \Delta \otimes 1)\Phi^{-1} \cdot (\Phi^{-1} \otimes 1) \cdot (\Delta \otimes 1 \otimes 1)\Phi \cdot (1 \otimes 1 \otimes \Delta)\Phi \\ &= \bar{X}_{\nu}\bar{X}_{\mu}X_{\rho}^{(1)}X_{\sigma} \otimes \bar{Y}_{\nu}^{(1)}\bar{Y}_{\mu}X_{\rho}^{(2)}Y_{\sigma} \otimes \bar{Y}_{\nu}^{(2)}\bar{Z}_{\mu}Y_{\rho}Z_{\sigma}^{(1)} \otimes \bar{Z}_{\nu}Z_{\rho}Z_{\sigma}^{(2)} \end{aligned} \quad (2.10)$$

where we have adopted the notation of equation (2.5) in (2.8) and the obvious notation in (2.9), (2.10) so that, for example

$$\Delta(X_{\nu}) = X_{\nu}^{(1)} \otimes X_{\nu}^{(2)}, \quad \text{etc.}$$

Equation (2.8) follows from applying $(1 \otimes m)(1 \otimes 1 \otimes \beta S)$ to equation (2.1), then using (2.7).

3 Uniqueness of the quasi-antipode.

For Hopf algebras the antipode S is uniquely determined as the inverse of the identity map on H under the convolution product. The quasi-antipode (S, α, β) for a QHA, is not unique. Nevertheless it is almost unique as the following result due to Drinfeld [6] (whose proof is similar to the one given below) shows:

Theorem 1. *Suppose H is also a QHA, but with quasi-antipode $(\tilde{S}, \tilde{\alpha}, \tilde{\beta})$ satisfying (2.6), (2.7). Then there exists a unique invertible $v \in H$ such that*

$$v\alpha = \tilde{\alpha}, \tilde{\beta}v = \beta, \tilde{S}(a) = vS(a)v^{-1}, \forall a \in H. \quad (3.11)$$

Explicitly

$$\begin{aligned} (i) \quad v &= \tilde{S}(X_\nu)\tilde{\alpha}Y_\nu\beta S(Z_\nu) = \tilde{S}(S^{-1}(\bar{X}_\nu))\tilde{S}(S^{-1}(\beta))\tilde{S}(\bar{Y}_\nu)\tilde{\alpha}\bar{Z}_\nu \\ (ii) \quad v^{-1} &= S(X_\nu)\alpha Y_\nu\tilde{\beta}\tilde{S}(Z_\nu) = \bar{X}_\nu\tilde{\beta}\tilde{S}(\bar{Y}_\nu)\tilde{S}(S^{-1}(\alpha))\tilde{S}(S^{-1}(\bar{Z}_\nu)) \end{aligned} \quad (3.12)$$

Proof. We proceed stepwise.

Applying $m \cdot (\tilde{S} \otimes 1)(1 \otimes \tilde{\alpha})$ to equation (2.8) gives

$$\tilde{S}(X_\nu a)\tilde{\alpha}Y_\nu\beta S(Z_\nu) = \tilde{S}(a_{(1)}^{(1)}X_\nu)\tilde{\alpha}a_{(1)}^{(2)}Y_\nu\beta S(Z_\nu)S(a_{(2)})$$

so that,

$$\tilde{S}(a)v = \tilde{S}(X_\nu)\tilde{S}(a_{(1)}^{(1)})\tilde{\alpha}a_{(1)}^{(2)}Y_\nu\beta S(Z_\nu)S(a_{(2)}) \stackrel{(2.7)}{=} vS(a), \quad \forall a \in H \quad (3.13)$$

where $m : H \otimes H \rightarrow H$ is the multiplication map $m(a \otimes b) = ab, \forall a, b \in H$.

Next observe, from equation (2.9) that, in view of (2.7),

$$\begin{aligned} v \otimes 1 &= \tilde{S}(X_\nu^{(1)}X_\mu\bar{X}_\rho)\tilde{\alpha}X_\nu^{(2)}Y_\mu\bar{X}_\sigma\bar{Y}_\rho^{(1)}\beta S(Y_\nu Z_\mu^{(1)}\bar{Y}_\sigma\bar{Y}_\rho^{(2)}) \otimes Z_\nu Z_\mu^{(2)}\bar{Z}_\sigma\bar{Z}_\rho \\ &= \tilde{S}(X_\mu)\tilde{\alpha}Y_\mu\bar{X}_\sigma\beta S(Z_\mu^{(1)}\bar{Y}_\sigma) \otimes Z_\mu^{(2)}\bar{Z}_\sigma. \end{aligned}$$

Applying $m \cdot (1 \otimes \alpha)$ from the left gives

$$\begin{aligned} v\alpha &= \tilde{S}(X_\mu)\tilde{\alpha}Y_\mu\bar{X}_\sigma\beta S(Z_\mu^{(1)}\bar{Y}_\sigma)\alpha Z_\mu^{(2)}\bar{Z}_\sigma \\ &= \tilde{\alpha}\bar{X}_\sigma\beta S(\bar{Y}_\sigma)\alpha\bar{Z}_\sigma \stackrel{(2.6)}{=} \tilde{\alpha}. \end{aligned} \quad (3.14)$$

From this it follows that

$$\begin{aligned} &\tilde{S}(S^{-1}(\bar{X}_\nu)) \cdot \tilde{S}(S^{-1}(\beta)) \cdot \tilde{S}(\bar{Y}_\nu)\tilde{\alpha}\bar{Z}_\nu \\ (3.14) \quad &\stackrel{=}{=} \tilde{S}(S^{-1}(\bar{X}_\nu)) \cdot \tilde{S}(S^{-1}(\beta))\tilde{S}(\bar{Y}_\nu) \cdot v\alpha\bar{Z}_\nu \\ (3.13) \quad &\stackrel{=}{=} v \cdot S(S^{-1}(\bar{X}_\nu)) \cdot S(S^{-1}(\beta)) \cdot S(\bar{Y}_\nu)\alpha\bar{Z}_\nu \\ &= v \cdot \bar{X}_\nu\beta S(\bar{Y}_\nu)\alpha\bar{Z}_\nu \stackrel{2.6}{=} v \end{aligned}$$

which proves (3.12)(i). To see v is invertible observe that

$$\begin{aligned} v \cdot S(X_\nu)\alpha Y_\nu\tilde{\beta}\tilde{S}(Z_\nu) &\stackrel{(3.13)}{=} \tilde{S}(X_\nu)v\alpha Y_\nu\tilde{\beta}\tilde{S}(Z_\nu) \\ &\stackrel{(3.14)}{=} \tilde{S}(X_\nu)\tilde{\alpha}Y_\nu\tilde{\beta}\tilde{S}(Z_\nu) \\ &\stackrel{(2.6)}{=} 1 \end{aligned}$$

so

$$v^{-1} = S(X_\nu)\alpha Y_\nu \tilde{\beta} \tilde{S}(Z_\nu)$$

as stated.

Now using equation (2.10) we have

$$\begin{aligned} 1 \otimes v^{-1} &= \bar{X}_\nu \bar{X}_\mu X_\rho^{(1)} X_\sigma \otimes S(\bar{Y}_\nu^{(1)} \bar{Y}_\mu X_\rho^{(2)} Y_\sigma) \alpha \bar{Y}_\nu^{(2)} \bar{Z}_\mu Y_\rho \bar{Z}_\sigma^{(1)} \tilde{\beta} \tilde{S}(\bar{Z}_\nu Z_\rho Z_\sigma^{(2)}) \\ &\stackrel{(2.7)}{=} \bar{X}_\mu X_\rho^{(1)} \otimes S(\bar{Y}_\mu X_\rho^{(2)}) \alpha \bar{Z}_\mu Y_\rho \tilde{\beta} \tilde{S}(Z_\rho). \end{aligned}$$

Applying $m \cdot (1 \otimes \beta)$ gives

$$\begin{aligned} \beta v^{-1} &= \bar{X}_\mu X_\rho^{(1)} \beta S(\bar{Y}_\mu X_\rho^{(2)}) \alpha \bar{Z}_\mu Y_\rho \tilde{\beta} \tilde{S}(Z_\rho) \\ &= \bar{X}_\mu \beta S(\bar{Y}_\mu) \alpha \bar{Z}_\mu \cdot \tilde{\beta} \stackrel{(2.6)}{=} \tilde{\beta} \end{aligned} \tag{3.15}$$

which completes the proof of (3.11). As to (3.12)(ii) observe that

$$\begin{aligned} &\bar{X}_\nu \tilde{\beta} \tilde{S}(\bar{Y}_\nu) \tilde{S}(S^{-1}(\alpha)) \tilde{S}(S^{-1}(\bar{Z}_\nu)) \\ &\stackrel{(3.15)}{=} \bar{X}_\nu \beta v^{-1} \tilde{S}(\bar{Y}_\nu) \tilde{S}(S^{-1}(\alpha)) \tilde{S}(S^{-1}(\bar{Z}_\nu)) \\ &\stackrel{(3.13)}{=} \bar{X}_\nu \beta S(\bar{Y}_\nu) S(S^{-1}(\alpha)) S(S^{-1}(\bar{Z}_\nu)) v^{-1} \\ &= \bar{X}_\nu \beta S(\bar{Y}_\nu) \alpha \bar{Z}_\nu \cdot v^{-1} \stackrel{(2.6)}{=} v^{-1} \end{aligned}$$

as required. It finally remains to prove uniqueness. Hence suppose $u \in H$ satisfies

$$uS(a) = \tilde{S}(a)u, \quad \forall a \in H, \quad u\alpha = \tilde{\alpha}, \quad \tilde{\beta}u = \beta.$$

Then

$$\begin{aligned} uv^{-1} &= u \cdot S(X_\nu) \alpha Y_\nu \tilde{\beta} \tilde{S}(Z_\nu) \\ &= \tilde{S}(X_\nu) u \alpha Y_\nu \tilde{\beta} \tilde{S}(Z_\nu) \\ &= \tilde{S}(X_\nu) \tilde{\alpha} Y_\nu \tilde{\beta} \tilde{S}(Z_\nu) \stackrel{(2.6)}{=} 1 \end{aligned}$$

which implies $u = v$ as required. $\square$

In the special case $\tilde{S} = S$ we obtain the following useful result.

Corollary. *Suppose H is also a QHA with quasi-antipode $(S, \tilde{\alpha}, \tilde{\beta})$. Then there is a unique invertible central element $v \in H$, given explicitly by equation (3.12)(i) (with $\tilde{S} = S$), such that*

$$v\alpha = \tilde{\alpha}, \quad \tilde{\beta}v = \beta.$$

It thus follows that the triple (S, α, β) satisfying (2.6), (2.7) for a QHA is not unique. Indeed following Theorem 1, for arbitrary invertible $v \in H$, the triple $(\tilde{S}, \tilde{\alpha}, \tilde{\beta})$ defined by

$$\tilde{S}(a) = vS(a)v^{-1}, \quad \forall a \in H; \quad \tilde{\alpha} = v\alpha, \quad \tilde{\beta} = \beta v^{-1}$$

is easily seen to satisfy (2.6), (2.7) and thus gives rise to a quasi-antipode $(\tilde{S}, \tilde{\alpha}, \tilde{\beta})$. Theorem 1 then shows that all such quasi-antipodes $(\tilde{S}, \tilde{\alpha}, \tilde{\beta})$ are obtainable this way: thus there is a 1–1 correspondence between the latter and invertible $v \in H$. We say that these structures are *equivalent* since they clearly give rise to equivalent QHA structures. Throughout we work with a fixed choice for the quasi-antipode (S, α, β) .

We conclude this section with the following useful result, proved in [13], concerning the opposite QHA structure on H :

Proposition 1. *H is also a QHA, with co-unit ϵ , under the opposite co-product and co-associator Δ^T , $\Phi^T \equiv \Phi_{321}^{-1}$ respectively, with quasi-antipode $(S^{-1}, \alpha^T = S^{-1}(\alpha), \beta^T = S^{-1}(\beta))$.*

The QHA $H^{cop} \equiv (H, \Delta^T, \epsilon, \Phi^T, S^{-1}, \alpha^T, \beta^T)$ is called the opposite QHA structure. We remark that above we have adopted the notation of [12] and [13] so that $\Delta^T = T \cdot \Delta$, T the usual twist map, and

$$\Phi_{321}^{-1} = \bar{Z}_\nu \otimes \bar{Y}_\nu \otimes \bar{X}_\nu.$$

This latter notation extends in a natural way and will be employed throughout.

4 Twisting

Let H be a quasi-bialgebra. Then $F \in H \otimes H$ is called a twist if it is invertible and satisfies the co-unit property

$$(\epsilon \otimes 1)F = (1 \otimes \epsilon)F = 1.$$

We recall that H is also a QBA with the same co-unit ϵ but with co-product and co-associator given by

$$\begin{aligned} \Delta_F(a) &= F\Delta(a)F^{-1}, \quad \forall a \in H \\ \Phi_F &= (F \otimes 1) \cdot (\Delta \otimes 1)F \cdot \Phi \cdot (1 \otimes \Delta)F^{-1} \cdot (1 \otimes F^{-1}), \end{aligned} \quad (4.16)$$

called the twisted structure induced by F . If moreover H is a QHA with quasi-antipode (S, α, β) then H is also a QHA under the above twisted structure with the *same* antipode S but with canonical elements

$$\alpha_F = m \cdot (1 \otimes \alpha)(S \otimes 1)F^{-1} \quad \beta_F = m \cdot (1 \otimes \beta)(1 \otimes S)F \quad (4.17)$$

respectively. A detailed proof of these well known results is given in [20]. We now investigate the behaviour of the operator v of Theorem 1 under the twisted structure induced by F .

4.1 Universality of v

Recall that the operator v is given by

$$v = \tilde{S}(X_\nu)\tilde{\alpha}Y_\nu\beta S(Z_\nu)$$

Let $F \in H \otimes H$ be an arbitrary twist. We use the following notation for the twist F and its inverse F^{-1} ,

$$F = f_i \otimes f^i, \quad F^{-1} = \bar{f}_i \otimes \bar{f}^i.$$

The twisted form of the co-associator is given by (4.16)

$$\Phi_F = X_\nu^F \otimes Y_\nu^F \otimes Z_\nu^F = f_i f_j^{(1)} X_\nu \bar{f}_k \otimes f^i f_j^{(2)} Y_\nu \bar{f}_{(1)}^k \bar{f}_l \otimes f^j Z_\nu \bar{f}_{(2)}^k \bar{f}^l. \quad (4.18)$$

For the twisted forms of the canonical elements we have from (4.17)

$$\begin{aligned} \tilde{\alpha}_F &= m \cdot (1 \otimes \tilde{\alpha})(\tilde{S} \otimes 1)F^{-1} = \tilde{S}(\bar{f}_p)\tilde{\alpha}\bar{f}^p \\ \beta_F &= m \cdot (1 \otimes \beta)(1 \otimes S)F = f_q \beta S(f^q). \end{aligned} \quad (4.19)$$

We note that

$$\tilde{S}(f_j)\tilde{\alpha}_F f^j \stackrel{(4.19)}{=} \tilde{S}(\bar{f}_p f_j)\tilde{\alpha}\bar{f}^p f^j = m \cdot (1 \otimes \alpha)(\tilde{S} \otimes 1)(F^{-1}F) = \tilde{\alpha} \quad (4.20)$$

and similarly,

$$\bar{f}_j \beta_F S(\bar{f}^j) = \beta. \quad (4.21)$$

The twisted form of v is given by

$$\begin{aligned} v_F &= \tilde{S}(X_\nu^F) \tilde{\alpha}_F Y_\nu^F \beta_F S(Z_\nu^F) \\ &\stackrel{(4.18)}{=} \tilde{S}(f_i f_j^{(1)} X_\nu \bar{f}_k) \tilde{\alpha}_F f^i f_j^{(2)} Y_\nu \bar{f}_{(1)}^k \bar{f}_l \beta_F S(f^j Z_\nu \bar{f}_{(2)}^k \bar{f}^l) \\ &= \tilde{S}(f_j^{(1)} X_\nu \bar{f}_k) \tilde{S}(f_i) \tilde{\alpha}_F f^i f_j^{(2)} Y_\nu \bar{f}_{(1)}^k \bar{f}_l \beta_F S(\bar{f}^l) S(f^j Z_\nu \bar{f}_{(2)}^k) \\ &\stackrel{(4.20)}{=} \tilde{S}(f_j^{(1)} X_\nu \bar{f}_k) \tilde{\alpha} f_j^{(2)} Y_\nu \bar{f}_{(1)}^k \bar{f}_l \beta_F S(\bar{f}^l) S(f^j Z_\nu \bar{f}_{(2)}^k) \\ &\stackrel{(4.21)}{=} \tilde{S}(f_j^{(1)} X_\nu \bar{f}_k) \tilde{\alpha} f_j^{(2)} Y_\nu \bar{f}_{(1)}^k \beta S(f^j Z_\nu \bar{f}_{(2)}^k) \\ &= \tilde{S}(X_\nu \bar{f}_k) \tilde{S}(f_j^{(1)}) \tilde{\alpha} f_j^{(2)} Y_\nu \bar{f}_{(1)}^k \beta S(\bar{f}_{(2)}^k) S(f^j Z_\nu) \\ &= \tilde{S}(X_\nu \bar{f}_k) \tilde{\alpha} Y_\nu \bar{f}_{(1)}^k \beta S(\bar{f}_{(2)}^k) S(Z_\nu) \\ &= \tilde{S}(X_\nu) \tilde{\alpha} Y_\nu \beta S(Z_\nu) = v, \end{aligned}$$

where, in the last two lines we have used the antipode properties of α, β (2.7) and the co-unit property of twists. We have thus proved

Theorem 2. *The operator v is universal (i.e. invariant under twisting).*

5 The Drinfeld twists

We turn our attention to the Drinfeld twist for the opposite structure of proposition 1. It is tempting to assume that F_δ^T qualifies as a Drinfeld twist for the opposite structure. However this is not true since the antipode for the latter is S^{-1} rather than S . We shall show that the Drinfeld twist for the opposite structure is in fact related to the second Drinfeld twist which we define below. We begin with a review of the Drinfeld twist.

5.1 The Drinfeld twist

Observe that Δ' defined by

$$\Delta'(a) = (S \otimes S) \Delta^T(S^{-1}(a)), \quad \forall a \in H \quad (5.22)$$

also determines a co-product on H . Associated with this co-product we have a new QHA structure on H , which was proved in [13] and which we restate here:

Proposition 2. *H is also a QHA with the same co-unit ϵ and antipode S but with co-product Δ' , co-associator $\Phi' = (S \otimes S \otimes S) \Phi_{321}$, and canonical elements $\alpha' = S(\beta)$, $\beta' = S(\alpha)$ respectively.*

Drinfeld has proved the remarkable result that this QHA structure is obtained by twisting with the Drinfeld twist, herein denoted F_δ , given explicitly by

$$\begin{aligned} (i) \quad F_\delta &= (S \otimes S) \Delta^T(X_\nu) \cdot \gamma \cdot \Delta(Y_\nu \beta S(Z_\nu)) \\ &= \Delta'(\bar{X}_\nu \beta S(\bar{Y}_\nu)) \cdot \gamma \cdot \Delta(\bar{Z}_\nu) \end{aligned}$$

where

$$(ii) \quad \gamma = S(B_i)\alpha C_i \otimes S(A_i)\alpha D_i$$

with

$$(iii) \quad A_i \otimes B_i \otimes C_i \otimes D_i = \begin{cases} (\Phi^{-1} \otimes 1) \cdot (\Delta \otimes 1 \otimes 1)\Phi \\ \text{or} \\ (1 \otimes \Phi) \cdot (1 \otimes 1 \otimes \Delta)\Phi^{-1} . \end{cases} \quad (5.23)$$

The inverse of F_δ is given explicitly by

$$(i) \quad F_\delta^{-1} = \Delta(\bar{X}_\nu) \cdot \bar{\gamma} \cdot \Delta'(S(\bar{Y}_\nu)\alpha\bar{Z}_\nu) \\ = \Delta(S(X_\nu)\alpha Y_\nu) \cdot \bar{\gamma} \cdot (S \otimes S)\Delta^T(Z_\nu)$$

where

$$(ii) \quad \bar{\gamma} = \bar{A}_i\beta S(\bar{D}_i) \otimes \bar{B}_i\beta S(\bar{C}_i)$$

with

$$(iii) \quad \bar{A}_i \otimes \bar{B}_i \otimes \bar{C}_i \otimes \bar{D}_i = \begin{cases} (\Delta \otimes 1 \otimes 1)\Phi^{-1} \cdot (\Phi \otimes 1) \\ \text{or} \\ (1 \otimes 1 \otimes \Delta)\Phi \cdot (1 \otimes \Phi^{-1}) . \end{cases} \quad (5.24)$$

The detailed proof that the QHA structure of proposition 2 is obtained by twisting with F_δ , as given in (5.23), and in particular

$$\Delta'(a) = F_\delta \Delta(a) F_\delta^{-1}, \quad \forall a \in H \quad (5.25)$$

is proved in [13]. We simply state here some properties of $\gamma, \bar{\gamma}$ proved in [13] and which are crucial to the demonstration of Drinfeld's result:

Proposition 3.

$$(i) \quad (S \otimes S)\Delta^T(a_{(1)}) \cdot \gamma \cdot \Delta(a_{(2)}) = \epsilon(a)\gamma, \quad \forall a \in H \\ (ii) \quad \Delta(a_{(1)}) \cdot \bar{\gamma} \cdot (S \otimes S)\Delta^T(a_{(2)}) = \epsilon(a)\bar{\gamma}, \quad \forall a \in H \\ (iii) \quad F_\delta \Delta(\alpha) = \gamma \quad , \quad \Delta(\beta)F_\delta^{-1} = \bar{\gamma} . \quad (5.26)$$

5.2 The second Drinfeld twist

Replacing S with S^{-1} we obtain yet another co-product Δ_0 on H :

$$\Delta_0(a) = (S^{-1} \otimes S^{-1})\Delta^T(S(a)), \quad \forall a \in H. \quad (5.22')$$

We have the following analogue of proposition 2, the proof of which parallels that of [13] proposition 4, but with S and S^{-1} interchanged:

Proposition 2' *H is also a QHA with the same co-unit ϵ and antipode S but with co-product Δ_0 , co-associator $\Phi_0 = (S^{-1} \otimes S^{-1} \otimes S^{-1})\Phi_{321}$ and canonical elements $\alpha_0 = S^{-1}(\beta)$, $\beta_0 = S^{-1}(\alpha)$ respectively.*

By symmetry we would expect this structure to be obtainable twisting. Indeed we have

Theorem 3. : The QHA structure of proposition 2' is obtained by twisting with

$$F_0 \equiv (S^{-1} \otimes S^{-1})F_\delta^T \quad (5.27)$$

herein referred to as the second Drinfeld twist, where F_δ is the Drinfeld twist and $F_\delta^T = T \cdot F_\delta$.

Proof. It is clear that F_0 is invertible with inverse $F_0^{-1} = (S^{-1} \otimes S^{-1})(F_\delta^T)^{-1}$ and qualifies as a twist. For the co-product we observe,

$$\begin{aligned} F_0 \Delta(a) F_0^{-1} &= (S^{-1} \otimes S^{-1}) F_\delta^T \cdot \Delta(a) \cdot (S^{-1} \otimes S^{-1}) (F_\delta^T)^{-1} \\ &= (S^{-1} \otimes S^{-1}) \cdot T \cdot [F_\delta^{-1} \cdot (S \otimes S) \Delta^T(a) \cdot F_\delta] \\ &= (S^{-1} \otimes S^{-1}) \cdot T \cdot [F_\delta^{-1} \Delta'(S(a)) F_\delta] \\ &\stackrel{(5.25)}{=} (S^{-1} \otimes S^{-1}) \cdot T \cdot \Delta(S(a)) = (S^{-1} \otimes S^{-1}) \Delta^T(S(a)) \\ &\stackrel{(5.22')}{=} \Delta_0(a), \quad \forall a \in H. \end{aligned}$$

The co-associator is slightly more complicated, though also simple. We have from Drinfeld's result

$$\Phi' \equiv (S \otimes S \otimes S) \Phi_{321} = (F_\delta \otimes 1) \cdot (\Delta \otimes 1) F_\delta \cdot \Phi \cdot (1 \otimes \Delta) F_\delta^{-1} \cdot (1 \otimes F_\delta^{-1})$$

which implies

$$\begin{aligned} (S \otimes S \otimes S) \Phi &= [(F_\delta \otimes 1) \cdot (\Delta \otimes 1) F_\delta \cdot \Phi \cdot (1 \otimes \Delta) F_\delta^{-1} \cdot (1 \otimes F_\delta^{-1})]_{321} \\ &= (1 \otimes F_\delta^T) \cdot (1 \otimes \Delta^T) F_\delta^T \cdot \Phi_{321} \cdot (\Delta^T \otimes 1) (F_\delta^T)^{-1} \cdot (F_\delta^{T-1} \otimes 1). \end{aligned}$$

Applying $(S^{-1} \otimes S^{-1} \otimes S^{-1})$ gives

$$\begin{aligned} \Phi &= (F_0^{-1} \otimes 1) \cdot (\Delta_0 \otimes 1) F_0^{-1} \cdot \Phi_0 \cdot (1 \otimes \Delta_0) F_0 \cdot (1 \otimes F_0) \\ &= (\Delta \otimes 1) F_0^{-1} \cdot (F_0^{-1} \otimes 1) \cdot \Phi_0 \cdot (1 \otimes F_0) \cdot (1 \otimes \Delta) F_0 \end{aligned}$$

with F_0 as in the Theorem. Thus

$$\Phi_0 = (F_0 \otimes 1) \cdot (\Delta \otimes 1) F_0 \cdot \Phi \cdot (1 \otimes \Delta) F_0^{-1} \cdot (1 \otimes F_0^{-1})$$

which shows that indeed Φ_0 is obtained from Φ by twisting with F_0 . The proof for the canonical elements is straightforward. $\square$

5.3 The Drinfeld twists for the opposite structure

Recall that under the opposite structure of proposition 1 H is a QHA with antipode S^{-1} , co-product Δ^T and co-associator $\Phi^T = \Phi_{321}^{-1}$. It follows that if F_δ^0 is the Drinfeld twist for this opposite structure then, $\forall a \in H$

$$\begin{aligned} F_\delta^0 \Delta^T(a) (F_\delta^0)^{-1} &= (\Delta^T)'(a) \\ &= (S^{-1} \otimes S^{-1}) \Delta(S(a)) = \Delta_0^T(a) \end{aligned}$$

since S^{-1} is the antipode for this structure. On the other hand if F_0 is the Drinfeld twist of equation (5.27) we have also

$$F_0^T \Delta^T(a) (F_0^T)^{-1} = \Delta_0^T(a)$$

with Δ_0 as in equation (5.22'). Here we show in fact that $F_\delta^0 = F_0^T$.

Before proceeding we note that the Drinfeld twist is given by the canonical expression of equation (5.23)(i) with γ as in (5.23)(ii) constructed from the operator of (5.23)(iii); viz

$$A_i \otimes B_i \otimes C_i \otimes D_i = \begin{cases} (\Phi^{-1} \otimes 1) \cdot (\Delta \otimes 1 \otimes 1) \Phi \\ \text{or} \\ (1 \otimes \Phi) \cdot (1 \otimes 1 \otimes \Delta) \Phi^{-1}. \end{cases}$$

This gives rise to two equivalent expansions for γ . Using the first expression we have, in obvious notation,

$$\begin{aligned} A_i \otimes B_i \otimes C_i \otimes D_i &= (\Phi^{-1} \otimes 1) \cdot (\Delta \otimes 1 \otimes 1) \Phi \\ &= \bar{X}_\nu X_\mu^{(1)} \otimes \bar{Y}_\nu X_\mu^{(2)} \otimes \bar{Z}_\nu Y_\mu \otimes Z_\mu \end{aligned}$$

which gives, upon substitution into (5.23)(ii),

$$\gamma = S(\bar{Y}_\nu X_\mu^{(2)}) \alpha \bar{Z}_\nu Y_\mu \otimes S(\bar{X}_\nu X_\mu^{(1)}) \alpha Z_\mu$$

which is the expression obtained in [13]. On the other hand using the second expression gives

$$\begin{aligned} A_i \otimes B_i \otimes C_i \otimes D_i &= (1 \otimes \Phi) \cdot (1 \otimes 1 \otimes \Delta) \Phi^{-1} \\ &= \bar{X}_\mu \otimes X_\nu \bar{Y}_\mu \otimes Y_\nu \bar{Z}_\mu^{(1)} \otimes Z_\nu \bar{Z}_\mu^{(2)} \end{aligned}$$

and substituting into (5.23)(ii) gives the alternative expansion

$$\gamma = S(X_\nu \bar{Y}_\mu) \alpha Y_\nu \bar{Z}_\mu^{(1)} \otimes S(\bar{X}_\mu) \alpha Z_\nu \bar{Z}_\mu^{(2)} \quad (5.28)$$

which is equivalent to the expression above [13].

Using (5.23)(i) for the opposite structure we have for the Drinfeld twist

$$F_\delta^0 = (S^{-1} \otimes S^{-1}) \Delta(X_\nu^0) \cdot \gamma^0 \cdot \Delta^T(Y_\nu^0 \beta^T S^{-1}(Z_\nu^0))$$

where we have used the fact that the co-product for the opposite structure is Δ^T , the antipode is S^{-1} , with canonical elements $\alpha^T = S^{-1}(\alpha)$, $\beta^T = S^{-1}(\beta)$ and where we have set

$$X_\nu^0 \otimes Y_\nu^0 \otimes Z_\nu^0 = \Phi^T = \Phi_{321}^{-1} \quad ,$$

which is the opposite co-associator, and where from (5.23)(ii)

$$\gamma^0 = S^{-1}(B_i^0) \alpha^T C_i^0 \otimes S^{-1}(A_i^0) \alpha^T D_i^0$$

with

$$\begin{aligned} A_i^0 \otimes B_i^0 \otimes C_i^0 \otimes D_i^0 &= [(\Phi^T)^{-1} \otimes 1] \cdot (\Delta^T \otimes 1 \otimes 1) \Phi^T \\ &= (\Phi_{321} \otimes 1) \cdot (\Delta^T \otimes 1 \otimes 1) \Phi_{321}^{-1}. \end{aligned}$$

In obvious notation the latter is given by

$$(\Phi_{321} \otimes 1) \cdot (\Delta^T \otimes 1 \otimes 1) \Phi_{321}^{-1} = Z_\nu \bar{Z}_\mu^{(2)} \otimes Y_\nu \bar{Z}_\mu^{(1)} \otimes X_\nu \bar{Y}_\mu \otimes \bar{X}_\mu$$

so that, using $\alpha^T = S^{-1}(\alpha)$,

$$\begin{aligned} \gamma^0 &= S^{-1}(Y_\nu \bar{Z}_\mu^{(1)}) S^{-1}(\alpha) X_\nu \bar{Y}_\mu \otimes S^{-1}(Z_\nu \bar{Z}_\mu^{(2)}) S^{-1}(\alpha) \bar{X}_\mu \\ &\stackrel{(5.28)}{=} (S^{-1} \otimes S^{-1})(\gamma). \end{aligned}$$

Thus we may write, using $\beta^T = S^{-1}(\beta)$,

$$F_\delta^0 = (S^{-1} \otimes S^{-1})\Delta(X_\nu^0) \cdot (S^{-1} \otimes S^{-1})\gamma \cdot \Delta^T(Y_\nu^0 S^{-1}(\beta) S^{-1}(Z_\nu^0))$$

so that, substituting

$$X_\nu^0 \otimes Y_\nu^0 \otimes Z_\nu^0 = \Phi^T = \Phi_{321}^{-1} = \bar{Z}_\nu \otimes \bar{Y}_\nu \otimes \bar{X}_\nu ,$$

gives

$$\begin{aligned} F_\delta^0 &= (S^{-1} \otimes S^{-1})\Delta(\bar{Z}_\nu) \cdot (S^{-1} \otimes S^{-1})\gamma \cdot \Delta^T(\bar{Y}_\nu S^{-1}(\beta) S^{-1}(\bar{X}_\nu)) \\ &= (S^{-1} \otimes S^{-1}) \cdot [(S \otimes S)\Delta^T(\bar{Y}_\nu S^{-1}(\bar{X}_\nu \beta)) \cdot \gamma \cdot \Delta(\bar{Z}_\nu)] \\ &= (S^{-1} \otimes S^{-1}) \cdot [\Delta'(\bar{X}_\nu \beta S(\bar{Y}_\nu)) \cdot \gamma \cdot \Delta(\bar{Z}_\nu)] \\ &\stackrel{(5.23)(i)}{=} (S^{-1} \otimes S^{-1})F_\delta \stackrel{(5.27)}{=} F_0^T. \end{aligned}$$

Thus we have proved

Proposition 4. *The Drinfeld twist for the opposite QHA structure of proposition 1 is given explicitly by*

$$F_\delta^0 = (S^{-1} \otimes S^{-1})F_\delta = F_0^T.$$

To see how F_δ^T fits into the picture we need to consider the second Drinfeld twist F_0 of Theorem 3 associated with the co-product of equation (5.22'). We have immediately from proposition 4

Corollary. *The second Drinfeld twist for the opposite structure is F_δ^T .*

Proof. Since the antipode for the opposite structure is S^{-1} , Theorem 3 implies that the second Drinfeld twist for this structure is $(S \otimes S)(F_\delta^0)^T$ where F_δ^0 is the Drinfeld twist for the opposite structure, given explicitly in proposition 4. It follows that the second Drinfeld twist for the opposite structure is

$$(S \otimes S) \cdot [(S^{-1} \otimes S^{-1})F_\delta^T] = F_\delta^T.$$

□

5.4 Twisting the Drinfeld twist

It is first useful to determine the behaviour of $\bar{\gamma}$ in equation (5.24)(ii) under an arbitrary twist $G \in H \otimes H$. Under the twisted structure induced by G the operator $\bar{\gamma}$ is twisted to $\bar{\gamma}_G$, given by equation (5.24)(ii,iii) for the twisted structure, so that

$$\begin{aligned} (i) \quad \bar{\gamma}_G &= \bar{A}_i^G \beta_G S(\bar{D}_i^G) \otimes \bar{B}_i^G \beta_G S(\bar{C}_i^G) \\ \text{where } (ii) \quad \bar{A}_i^G \otimes \bar{B}_i^G \otimes \bar{C}_i^G \otimes \bar{D}_i^G &= (\Delta_G \otimes 1 \otimes 1) \Phi_G^{-1} \cdot (\Phi_G \otimes 1). \end{aligned} \tag{5.29}$$

We have

Proposition 5. *Let $G = g_i \otimes g^i \in H \otimes H$ be a twist on a QHA H . Then*

$$\bar{\gamma}_G = G \cdot \Delta(g_i) \cdot \bar{\gamma} \cdot (S \otimes S)(G^T \Delta^T(g^i)).$$

Proof. Throughout we write

$$G^{-1} = \bar{g}_i \otimes \bar{g}^i.$$

For the RHS of equation (5.29)(ii) we have

$$\begin{aligned} (\Delta_G \otimes 1 \otimes 1) \Phi_G^{-1} \cdot (\Phi_G \otimes 1) &= (\Delta_G \otimes 1 \otimes 1) \cdot \\ &[(1 \otimes G) \cdot (1 \otimes \Delta) G \cdot \Phi^{-1} \cdot (\Delta \otimes 1) G^{-1} \cdot (G^{-1} \otimes 1)] \cdot \\ &\{[(G \otimes 1) \cdot (\Delta \otimes 1) G \cdot \Phi \cdot (1 \otimes \Delta) G^{-1} \cdot (1 \otimes G^{-1})] \otimes 1\} \end{aligned}$$

where we have used equation (4.16) for Φ_G and its inverse, thus

$$\begin{aligned} (\Delta_G \otimes 1 \otimes 1) \Phi_G^{-1} \cdot (\Phi_G \otimes 1) &= \\ (1 \otimes 1 \otimes G) \cdot (\Delta_G \otimes \Delta) G \cdot (\Delta_G \otimes 1 \otimes 1) \Phi^{-1} \cdot [(\Delta_G \otimes 1) \Delta \otimes 1] G^{-1} \\ &\cdot [(\Delta_G \otimes 1) G^{-1} \otimes 1] \cdot (G \otimes 1 \otimes 1) \cdot [(\Delta \otimes 1) G \otimes 1] \cdot (\Phi \otimes 1) \\ &\cdot [(1 \otimes \Delta) G^{-1} \otimes 1] \cdot (1 \otimes G^{-1} \otimes 1) \\ &= (G \otimes G) \cdot (\Delta \otimes \Delta) G \cdot (\Delta \otimes 1 \otimes 1) \Phi^{-1} \cdot [(\Delta \otimes 1) \Delta \otimes 1] G^{-1} \\ &\cdot [(\Delta \otimes 1) G^{-1} \otimes 1] \cdot [(\Delta \otimes 1) G \otimes 1] \cdot (\Phi \otimes 1) \\ &\cdot [(1 \otimes \Delta) G^{-1} \otimes 1] \cdot (1 \otimes G^{-1} \otimes 1) \\ &= (G \otimes G) \cdot (\Delta \otimes \Delta) G \cdot (\Delta \otimes 1 \otimes 1) \Phi^{-1} \cdot [(\Delta \otimes 1) \Delta \otimes 1] G^{-1} \\ &\cdot (\Phi \otimes 1) \cdot [(1 \otimes \Delta) G^{-1} \otimes 1] \cdot (1 \otimes G^{-1} \otimes 1) \\ &\stackrel{(2.1)}{=} (G \otimes G) \cdot (\Delta \otimes \Delta) G \cdot \{(\Delta \otimes 1 \otimes 1) \Phi^{-1} \cdot (\Phi \otimes 1)\} \\ &\cdot [(1 \otimes \Delta) \Delta \otimes 1] G^{-1} \cdot [(1 \otimes \Delta) G^{-1} \otimes 1] \cdot (1 \otimes G^{-1} \otimes 1). \end{aligned}$$

Now using the notation of equation (5.24)(iii) we have

$$(\Delta \otimes 1 \otimes 1) \Phi^{-1} \cdot (\Phi \otimes 1) = \bar{A}_i \otimes \bar{B}_i \otimes \bar{C}_i \otimes \bar{D}_i$$

so that in the notation of equation (5.29)(i)

$$\begin{aligned} \bar{A}_i^G \otimes \bar{B}_i^G \otimes \bar{C}_i^G \otimes \bar{D}_i^G &= (\Delta_G \otimes 1 \otimes 1) \Phi_G^{-1} \cdot (\Phi_G \otimes 1) \\ &= (G \otimes G) \cdot (\Delta \otimes \Delta) G \cdot \{\bar{A}_i \otimes \bar{B}_i \otimes \bar{C}_i \otimes \bar{D}_i\} \\ &\cdot [(1 \otimes \Delta) \Delta \otimes 1] G^{-1} \cdot [(1 \otimes \Delta) G^{-1} \otimes 1] \cdot (1 \otimes G^{-1} \otimes 1) \\ &= g_s g_j^{(1)} \bar{A}_i \bar{g}_l^{(1)} \bar{g}_k \otimes g^s g_j^{(2)} \bar{B}_i \bar{g}_{l(1)}^{(2)} \bar{g}_{(1)}^k \bar{g}_m \otimes g_t g_{(1)}^j \bar{C}_i \bar{g}_{l(2)}^{(2)} \bar{g}_{(2)}^k \bar{g}^m \otimes g^t g_{(2)}^j \bar{D}_i \bar{g}^l \end{aligned}$$

where we have used the obvious notation, so that

$$\begin{aligned} \Delta(g_i) &= g_i^{(1)} \otimes g_i^{(2)}, \\ (1 \otimes \Delta) \Delta(g_i) &= g_i^{(1)} \otimes \Delta(g_i^{(2)}) = g_i^{(1)} \otimes g_{i(1)}^{(2)} \otimes g_{i(2)}^{(2)}, \text{ etc} \end{aligned}$$

and all repeated indices are understood to be summed over.

Substituting into equation (5.29)(i) gives

$$\begin{aligned} \bar{\gamma}_G &= g_s g_j^{(1)} \bar{A}_i \bar{g}_l^{(1)} \bar{g}_k \beta_G S(g^t g_{(2)}^j \bar{D}_i \bar{g}^l) \\ &\otimes g^s g_j^{(2)} \bar{B}_i \bar{g}_{l(1)}^{(2)} \bar{g}_{(1)}^k \bar{g}_m \beta_G S(g_t g_{(1)}^j \bar{C}_i \bar{g}_{l(2)}^{(2)} \bar{g}_{(2)}^k \bar{g}^m) \\ &= g_s g_j^{(1)} \bar{A}_i \bar{g}_l^{(1)} \bar{g}_k \beta_G S(g^t g_{(2)}^j \bar{D}_i \bar{g}^l) \\ &\otimes g^s g_j^{(2)} \bar{B}_i \bar{g}_{l(1)}^{(2)} \bar{g}_{(1)}^k \bar{g}_m \beta_G S(\bar{g}^m) S(\bar{g}_{(2)}^k) S(\bar{g}_{l(2)}^{(2)}) S(g_t g_{(1)}^j \bar{C}_i). \end{aligned}$$

Now using

$$\bar{g}_m \beta_G S(\bar{g}^m) = (\beta_G)_{G^{-1}} = \beta_{G^{-1}G} = \beta \quad (5.30)$$

and making repeated use of equation (2.7) gives

$$\begin{aligned} \bar{\gamma}_G &= g_s g_j^{(1)} \bar{A}_i \bar{g}_l^{(1)} \bar{g}_k \beta_G S(g^t g_{(2)}^j \bar{D}_i \bar{g}^l) \\ &\quad \otimes g^s g_j^{(2)} \bar{B}_i \bar{g}_{l(1)}^{(2)} \bar{g}_{(1)}^k \beta S(\bar{g}_{(2)}^k) S(\bar{g}_{l(2)}^{(2)}) S(g_t g_{(1)}^j \bar{C}_i) \\ &= g_s g_j^{(1)} \bar{A}_i \bar{g}_l \beta_G S(\bar{g}^l) S(g^t g_{(2)}^j \bar{D}_i) \otimes g^s g_j^{(2)} \bar{B}_i \beta S(g_t g_{(1)}^j \bar{C}_i) \\ &\stackrel{(5.30)}{=} g_s g_j^{(1)} \bar{A}_i \beta S(\bar{D}_i) S(g^t g_{(2)}^j) \otimes g^s g_j^{(2)} \bar{B}_i \beta S(\bar{C}_i) S(g_t g_{(1)}^j) \\ &\stackrel{(5.24)(ii)}{=} (g_s g_j^{(1)} \otimes g^s g_j^{(2)}) \cdot \bar{\gamma} \cdot (S \otimes S)(g^t g_{(2)}^j \otimes g_t g_{(1)}^j) \\ &= G \cdot \Delta(g_j) \cdot \bar{\gamma} \cdot (S \otimes S)(G^T \cdot \Delta^T(g^j)) \end{aligned}$$

which proves the result. $\square$

We are now in a position to determine the action of an arbitrary twist $G \in H \otimes H$ on the inverse Drinfeld twist F_δ^{-1} , given in equation (5.24). Under the twisted structure induced by G , F_δ^{-1} is twisted to $(F_\delta^G)^{-1} \equiv (F_\delta^{-1})_G$, given as in equation (5.24), but in terms of the twisted structure, so that, with the notation of equation (5.29), we have from (5.24)(i)

$$(F_\delta^G)^{-1} = \Delta_G(S(X_\nu^G) \alpha_G Y_\nu^G) \cdot \bar{\gamma}_G \cdot (S \otimes S) \Delta_G^T(Z_\nu^G)$$

with $\bar{\gamma}_G$ as in proposition 5.

In obvious notation we may write

$$\begin{aligned} X_\nu^G \otimes Y_\nu^G \otimes Z_\nu^G = \Phi_G &= (G \otimes 1) \cdot (\Delta \otimes 1) G \cdot \Phi \cdot (1 \otimes \Delta) G^{-1} \cdot (1 \otimes G^{-1}) \\ &= g_i g_j^{(1)} X_\nu \bar{g}_k \otimes g^i g_j^{(2)} Y_\nu \bar{g}_{(1)}^k \bar{g}_l \otimes g^j Z_\nu \bar{g}_{(2)}^k \bar{g}^l \end{aligned}$$

which implies

$$\begin{aligned} (F_\delta^G)^{-1} &= \Delta_G[S(g_i g_j^{(1)} X_\nu \bar{g}_k) \alpha_G g^i g_j^{(2)} Y_\nu \bar{g}_{(1)}^k \bar{g}_l] \cdot \bar{\gamma}_G \cdot (S \otimes S) \Delta_G^T(g^j Z_\nu \bar{g}_{(2)}^k \bar{g}^l) \\ &= \Delta_G[S(X_\nu \bar{g}_k) S(g_j^{(1)}) S(g_i) \alpha_G g^i g_j^{(2)} Y_\nu \bar{g}_{(1)}^k \bar{g}_l] \cdot \bar{\gamma}_G \\ &\quad \cdot (S \otimes S) \Delta_G^T(g^j Z_\nu \bar{g}_{(2)}^k \bar{g}^l). \end{aligned}$$

Using

$$S(g_i) \alpha_G g^i = (\alpha_G)_{G^{-1}} = \alpha_{G^{-1}G} = \alpha,$$

and equation (2.7), then gives

$$\begin{aligned}
(F_\delta^G)^{-1} &= \Delta_G[S(X_\nu \bar{g}_k) \alpha Y_\nu \bar{g}_{(1)}^k \bar{g}_l] \cdot \bar{\gamma}_G \cdot (S \otimes S) \Delta_G^T(Z_\nu \bar{g}_{(2)}^k \bar{g}^l) \\
&= G \cdot \Delta[S(X_\nu \bar{g}_k) \alpha Y_\nu \bar{g}_{(1)}^k \bar{g}_l] \cdot G^{-1} \cdot \bar{\gamma}_G \\
&\quad \cdot (S \otimes S)(G^T)^{-1} \cdot (S \otimes S) \Delta^T(Z_\nu \bar{g}_{(2)}^k \bar{g}^l) \cdot (S \otimes S) G^T \\
&\stackrel{\text{prop. (5)}}{=} G \cdot \Delta[S(X_\nu \bar{g}_k) \alpha Y_\nu \bar{g}_{(1)}^k \bar{g}_l] \cdot \Delta(g_i) \cdot \bar{\gamma} \\
&\quad \cdot (S \otimes S) \Delta^T(g^i) \cdot (S \otimes S) \Delta^T(Z_\nu \bar{g}_{(2)}^k \bar{g}^l) \cdot (S \otimes S) G^T \\
&= G \cdot \Delta[S(X_\nu \bar{g}_k) \alpha Y_\nu \bar{g}_{(1)}^k] \cdot \Delta(\bar{g}_l) \Delta(g_i) \cdot \bar{\gamma} \\
&\quad \cdot (S \otimes S) \Delta^T(g^i) \cdot (S \otimes S) \Delta^T(\bar{g}^l) \cdot (S \otimes S) \Delta^T(Z_\nu \bar{g}_{(2)}^k) \\
&\quad \cdot (S \otimes S) G^T \\
&= G \cdot \Delta[S(X_\nu \bar{g}_k) \alpha Y_\nu \bar{g}_{(1)}^k] \cdot \Delta(\bar{g}_l g_i) \cdot \gamma \\
&\quad \cdot (S \otimes S) \Delta^T(\bar{g}^l g^i) \cdot (S \otimes S) \Delta^T(Z_\nu \bar{g}_{(2)}^k) \cdot (S \otimes S) G^T \\
&= G \cdot \Delta[S(X_\nu \bar{g}_k) \alpha Y_\nu] \cdot \Delta(\bar{g}_{(1)}^k) \cdot \gamma \\
&\quad \cdot (S \otimes S) \Delta^T(\bar{g}_{(2)}^k) \cdot (S \otimes S) \Delta^T(Z_\nu) \cdot (S \otimes S) G^T
\end{aligned}$$

where we have used the obvious result that

$$\bar{g}_l g_i \otimes \bar{g}^l g^i = G^{-1} G = 1 \otimes 1.$$

It then follows from proposition 3 that

$$\begin{aligned}
(F_\delta^G)^{-1} &= G \cdot \Delta[S(X_\nu) \alpha Y_\nu] \cdot \bar{\gamma} \cdot (S \otimes S) \Delta^T(Z_\nu) \cdot (S \otimes S) G^T \\
&\stackrel{(5.24)(i)}{=} G \cdot F_\delta^{-1} \cdot (S \otimes S) G^T.
\end{aligned}$$

We have thus proved

Theorem 4. *Let $G \in H \otimes H$ be a twist on a QHA H . Then under the twisted structure induced by G , F_δ^{-1} is twisted to*

$$(F_\delta^G)^{-1} \equiv (F_\delta^{-1})_G = G \cdot F_\delta^{-1} \cdot (S \otimes S) G^T.$$

Equivalently, the Drinfeld twist is twisted to

$$F_\delta^G \equiv (F_\delta)_G = (S \otimes S)(G^T)^{-1} \cdot F_\delta \cdot G^{-1}.$$

Corollary. F_0 as in equation (5.27) is twisted to

$$F_0^G \equiv (F_0)_G = (S^{-1} \otimes S^{-1})(G^T)^{-1} \cdot F_0 \cdot G^{-1}.$$

Proof. Follows from the definition of $F_0 \equiv (S^{-1} \otimes S^{-1}) F_\delta^T$ and the Theorem above. $\square$

When H is quasi-triangular the opposite structure of proposition 1 is obtainable, up to equivalence modulo (S, α, β) , via twisting. In such a case the results of Section 3 have further useful consequences.

6 Quasi-triangular QHAs

A QHA H is called quasi-triangular if there exists an invertible element

$$\mathcal{R} = \sum_i e_i \otimes e^i \in H \otimes H$$

called the R -matrix, such that

$$\begin{aligned} (i) \quad & \Delta^T(a)\mathcal{R} = \mathcal{R}\Delta(a), \quad \forall a \in H \\ (ii) \quad & (\Delta \otimes 1)\mathcal{R} = \Phi_{231}^{-1}\mathcal{R}_{13}\Phi_{132}\mathcal{R}_{23}\Phi_{123}^{-1} \\ (iii) \quad & (1 \otimes \Delta)\mathcal{R} = \Phi_{312}\mathcal{R}_{13}\Phi_{213}^{-1}\mathcal{R}_{12}\Phi_{123}, \end{aligned} \tag{6.31}$$

where

$$\mathcal{R}_{12} = e_i \otimes e^i \otimes 1, \quad \mathcal{R}_{13} = e_i \otimes 1 \otimes e^i, \quad \text{etcetera.}$$

We first summarise some well known results for quasi-triangular QHAs. It was shown in [13] that

Proposition 1' *Under the opposite QHA structure of proposition 1, H is also quasi-triangular with R -matrix $\mathcal{R}^T = T \cdot \mathcal{R}$, called the opposite R -matrix.*

It follows from (6.31) (ii,iii) that

$$(\epsilon \otimes 1)\mathcal{R} = (1 \otimes \epsilon)\mathcal{R} = 1$$

so that $\mathcal{R}$ qualifies as a twist. Moreover if $F \in H \otimes H$ is any twist then, as shown in [13], H is also quasi-triangular under the twisted structure of equations (4.16, 4.17) with R -matrix

$$\mathcal{R}_F = F^T \mathcal{R} F^{-1}. \tag{6.32}$$

It was shown in [13] that

Proposition 6. *Under the QHA of proposition 2, H is also quasi-triangular with R -matrix*

$$\mathcal{R}' = (S \otimes S)\mathcal{R}.$$

We have seen that the QHA structure of proposition 2 is obtainable by twisting with the Drinfeld twist F_δ . It was further shown in [13] that the full structure of proposition 6 is also obtained by twisting with F_δ which, in view of equation (6.32), is equivalent to

$$(S \otimes S)\mathcal{R} = F_\delta^T \mathcal{R} F_\delta^{-1}. \tag{6.33}$$

This result in fact follows from the following relation

$$(S \otimes S)\mathcal{R} \cdot \gamma = \gamma^T \mathcal{R},$$

where $\gamma^T = T \cdot \gamma$, proved in [13]. In view of proposition 3 this last equation is equivalent to

$$\mathcal{R}\bar{\gamma} = \bar{\gamma}^T \cdot (S \otimes S)\mathcal{R}$$

where $\bar{\gamma}^T = T \cdot \bar{\gamma}$, with γ and $\bar{\gamma}$ as in equations (5.23, 5.24).

In view of (6.31) (i) the opposite co-product is obtained from Δ by twisting with $\mathcal{R}$. In fact we have the following result proved in [13] :

Proposition 7. *The opposite structure of propositions 1, 1' is obtainable by twisting with the R -matrix $\mathcal{R}$ but with antipode S and canonical elements $\alpha_{\mathcal{R}}, \beta_{\mathcal{R}}$ respectively.*

Above $\alpha_{\mathcal{R}}, \beta_{\mathcal{R}}$ are given by equation (4.17), so that

$$(i) \quad \alpha_{\mathcal{R}} = m \cdot (1 \otimes \alpha)(S \otimes 1)\mathcal{R}^{-1}, \quad \beta_{\mathcal{R}} = m \cdot (1 \otimes \beta)(1 \otimes S)\mathcal{R}.$$

Below we set

$$(ii) \quad \mathcal{R} = e_i \otimes e^i, \quad \mathcal{R}^{-1} = \bar{e}_i \otimes \bar{e}^i$$

in terms of which we may write

$$(iii) \quad \alpha_{\mathcal{R}} = S(\bar{e}_i)\alpha\bar{e}^i, \quad \beta_{\mathcal{R}} = e_i\beta S(e^i). \quad (6.34)$$

Thus with the co-product Δ^T and co-associator $\Phi^T = \Phi_{321}^{-1}$ of proposition 1 we have two QHA structures with differing quasi-antipodes $(S, \alpha_{\mathcal{R}}, \beta_{\mathcal{R}})$ and $(S^{-1}, \alpha^T, \beta^T)$ where, from proposition 1, $\alpha^T = S^{-1}(\alpha)$, $\beta^T = S^{-1}(\beta)$. It follows from Theorem 1 that

Theorem 5. *There exists a unique invertible $u \in H$ such that*

$$S(a) = uS^{-1}(a)u^{-1}, \text{ or } S^2(a) = uau^{-1}, \quad \forall a \in H$$

and

$$uS^{-1}(\alpha) = \alpha_{\mathcal{R}}, \quad \beta_{\mathcal{R}}u = S^{-1}(\beta). \quad (6.35)$$

Explicitly,

$$\begin{aligned} u &= S(Y_\nu \beta S(Z_\nu))\alpha_{\mathcal{R}}X_\nu = S(\bar{Z}_\nu)\alpha_{\mathcal{R}}\bar{Y}_\nu S^{-1}(\beta)S^{-1}(\bar{X}_\nu) \\ u^{-1} &= Z_\nu \beta_{\mathcal{R}}S(S(X_\nu)\alpha Y_\nu) = S^{-1}(\bar{Z}_\nu)S^{-1}(\alpha)\bar{Y}_\nu \beta_{\mathcal{R}}S(\bar{X}_\nu). \end{aligned} \quad (6.36)$$

Above we have used the fact that the opposite QHA structure has co-associator $\Phi^T = \Phi_{321}^{-1}$ and quasi-antipode $(S^{-1}, \alpha^T, \beta^T)$. We have then applied Theorem 1 with $(\tilde{S}, \tilde{\alpha}, \tilde{\beta}) = (S, \alpha_{\mathcal{R}}, \beta_{\mathcal{R}})$ to give the result.

The above gives the u -operator of Drinfeld-Reshetikhin [5, 18]. It differs from, but is related to, the u -operator of Altschuler and Coste [1]. To see how the latter arises, it is easily seen that $\tilde{\mathcal{R}} \equiv (\mathcal{R}^T)^{-1}$ also satisfies equation (6.31) and thus constitutes an R -matrix. Thus proposition 7 and Theorem 3 also hold with $\mathcal{R}$ replaced by $\tilde{\mathcal{R}}$. This implies the existence of a unique invertible $\tilde{u} \in H$ such that

$$S^2(a) = \tilde{u}a\tilde{u}^{-1}, \quad \forall a \in H$$

and

$$\tilde{u}S^{-1}(\alpha) = \alpha_{\tilde{\mathcal{R}}}, \quad \beta_{\tilde{\mathcal{R}}}\tilde{u} = S^{-1}(\beta)$$

with $\alpha_{\tilde{\mathcal{R}}}, \beta_{\tilde{\mathcal{R}}}$ as in equation (6.34) but with $\mathcal{R}$ replaced by $\tilde{\mathcal{R}}$. Explicitly we have, in this case,

$$\begin{aligned} \tilde{u} &= S(Y_\nu \beta S(Z_\nu))\alpha_{\tilde{\mathcal{R}}}X_\nu = S(\bar{Z}_\nu)\alpha_{\tilde{\mathcal{R}}}\bar{Y}_\nu S^{-1}(\beta)S^{-1}(\bar{X}_\nu) \\ \tilde{u}^{-1} &= Z_\nu \beta_{\tilde{\mathcal{R}}}S(S(X_\nu)\alpha Y_\nu) = S^{-1}(\bar{Z}_\nu)S^{-1}(\alpha)\bar{Y}_\nu \beta_{\tilde{\mathcal{R}}}S(\bar{X}_\nu). \end{aligned} \quad (6.37)$$

Then, as can be seen from [12] $\tilde{u}$ is precisely the u -operator of Altschuler and Coste.

To see the relation between u and $\tilde{u}$ we first note that $uS(u) = S(u)u$ is central. This follows by applying S to $S(a) = uS^{-1}(a)u^{-1}$, giving

$$S^2(a) = S(u^{-1})aS(u), \quad \forall a \in H.$$

Before proceeding it is worth noting the following

Lemma 1.

$$\begin{aligned} (i) \quad & \beta_{\tilde{\mathcal{R}}} = S(u)S(\beta), \quad \alpha_{\tilde{\mathcal{R}}} = S(\alpha)S(u^{-1}) \\ (ii) \quad & \beta_{\mathcal{R}} = S(\tilde{u})S(\beta), \quad \alpha_{\mathcal{R}} = S(\alpha)S(\tilde{u}^{-1}). \end{aligned} \tag{6.38}$$

Proof. By symmetry it suffices to prove (i). Now

$$\begin{aligned} \beta_{\tilde{\mathcal{R}}} &= m \cdot (1 \otimes \beta)(1 \otimes S)(\mathcal{R}^T)^{-1} = \bar{e}^i \beta S(\bar{e}_i) \\ &\stackrel{(6.35)}{=} \bar{e}^i S(\beta_{\mathcal{R}} u) S(\bar{e}_i) = \bar{e}^i S(u) S(\beta_{\mathcal{R}}) S(\bar{e}_i) \\ &= \bar{e}^i S(u) S[e_j \beta S(e^j)] S(\bar{e}_i) \\ &= \bar{e}^i S(u) S^2(e^j) S(\beta) S(e_j) S(\bar{e}_i) \\ &= S(u) S^2(\bar{e}^i) S^2(e^j) S(\beta) S(e_j) S(\bar{e}_i) \\ &= S(u) S^2(\bar{e}^i e^j) S(\beta) S(\bar{e}_i e_j) = S(u) S(\beta) \end{aligned}$$

where we have used the obvious result

$$\bar{e}_i e_j \otimes \bar{e}^i e^j = \mathcal{R}^{-1} \mathcal{R} = 1 \otimes 1.$$

Similarly

$$\begin{aligned} \alpha_{\tilde{\mathcal{R}}} &= m \cdot (1 \otimes \alpha)(S \otimes 1) R^T = S(e^i) \alpha e_i \\ &\stackrel{(6.35)}{=} S(e^i) S(u^{-1} \alpha_{\mathcal{R}}) e_i = S(e^i) S(\alpha_{\mathcal{R}}) S(u^{-1}) e_i \\ &= S(e^i) S[S(\bar{e}_j) \alpha \bar{e}^j] S(u^{-1}) e_i \\ &= S(e^i) S(\bar{e}^j) S(\alpha) S^2(\bar{e}_j) S(u^{-1}) e_i \\ &= S(e^i) S(\bar{e}^j) S(\alpha) S^2(\bar{e}_j) S^2(e_i) S(u^{-1}) \\ &= S(\bar{e}^j e^i) S(\alpha) S^2(\bar{e}_j e_i) S(u^{-1}) = S(\alpha) S(u^{-1}). \end{aligned}$$

□

We are now in a position to prove

Lemma 2.

$$\tilde{u} = S(u^{-1})$$

Proof. From equation (6.37) we have

$$\begin{aligned} \tilde{u} &= S(Y_\nu \beta S(Z_\nu)) \alpha_{\tilde{\mathcal{R}}} X_\nu \\ &\stackrel{(6.38)(i)}{=} S(Y_\nu \beta S(Z_\nu)) S(\alpha) S(u^{-1}) X_\nu \\ &= S(Y_\nu \beta S(Z_\nu)) S(\alpha) S^2(X_\nu) S(u^{-1}) \\ &= S[S(X_\nu) \alpha Y_\nu \beta S(Z_\nu)] S(u^{-1}) \\ &\stackrel{(2.6)}{=} S(u^{-1}). \end{aligned}$$

□

The above result clearly shows the connection between the u -operator of Theorem 3 and that due to Altschuler and Coste. Obviously the existence of the u -operator in the quasi-triangular case is a direct consequence of Theorem 1 and proposition 7, the latter showing the equivalence

of the opposite structure of proposition 1 with that due to twisting with $\mathcal{R}$. In the case H is not quasi-triangular, this opposite structure is not in general obtainable by a twist.

The operators u and $\tilde{u}$ are special cases of the v operator of Theorem 1, it follows then from Theorem 2, that

Theorem 6. *The operators u and $\tilde{u}$ are invariant under twisting.*

In section 3 we discussed the uniqueness of the quasi-antipode (S, α, β) , but nothing has been said about the uniqueness of the twisted structures or the R -matrix in the quasi-triangular case. This is intimately connected with the quasi-cocycle condition to which we now turn.

7 The quasi-cocycle condition

The set of twists on a QHA H forms a group, moreover, the twisted structure of equations (4.16, 4.17) induced on a QHA H preserves this group structure in the following sense.

Lemma 3. *Let $F, G \in H \otimes H$ be twists on a QHA H . Then in the notation of equations (4.16, 4.17)*

$$\begin{aligned} (i) \quad & \Delta_{FG} = (\Delta_G)_F, \quad \Phi_{FG} = (\Phi_G)_F \\ (ii) \quad & \alpha_{FG} = (\alpha_G)_F, \quad \beta_{FG} = (\beta_G)_F. \end{aligned}$$

Moreover, if H is quasi-triangular then

$$(iii) \quad \mathcal{R}_{FG} = (\mathcal{R}_G)_F. \quad (7.39)$$

In other words the structure obtained from twisting with G and then with F is the same as twisting with the twist FG . It is important that the right hand side of equation (7.39) is interpreted correctly, e.g. $(\Phi_G)_F$ is given as in equation (4.16) but with Φ replaced by Φ_G and Δ by Δ_G etc.

Given any QBA H we may impose on a twist $F \in H \otimes H$ the following condition

$$(F \otimes 1) \cdot (\Delta \otimes 1)F \cdot \Phi = \Phi \cdot (1 \otimes F) \cdot (1 \otimes \Delta)F \quad (7.40)$$

which we call the *quasi-cocycle condition*.

When $\Phi = 1 \otimes 1 \otimes 1$ this reduces to the usual cocycle condition on Hopf algebras. In the notation of equation (4.16), the quasi-cocycle condition is equivalent to

$$\Phi_F = \Phi. \quad (7.40')$$

Thus twisting on a QBA by a twist F satisfying the quasi-cocycle condition results in a QBA structure with the same co-associator.

It is thus not surprising that the quasi-cocycle condition (7.40) is intimately related to the uniqueness of twisted structures on a QHA H . Indeed, if $F, G \in H \otimes H$ are twists giving rise to the *same* QBA structure, so that

$$\Delta_F = \Delta_G, \quad \Phi_F = \Phi_G \quad (7.41)$$

then $C \equiv F^{-1}G$ must commute with the co-product Δ and satisfy the quasi-cocycle condition. Indeed in view of lemma 3 we have

$$\begin{aligned} \Delta_C &= \Delta_{F^{-1}G} = (\Delta_G)_{F^{-1}} \stackrel{(7.41)}{=} (\Delta_F)_{F^{-1}} = \Delta_{F^{-1}F} = \Delta \\ \Phi_C &= \Phi_{F^{-1}G} = (\Phi_G)_{F^{-1}} \stackrel{(7.41)}{=} (\Phi_F)_{F^{-1}} = \Phi_{F^{-1}F} = \Phi. \end{aligned}$$

This leads to the following

Definition 3. A twist $C \in H \otimes H$ on any QBA H is called compatible if

- (i) C commutes with the co-product Δ
- (ii) C satisfies the quasi-cocycle condition .

In other words twisting a QBA H with a compatible twist C gives exactly the same QBA structure. The set of compatible twists on H thus forms a subgroup of the group of twists on H .

Proposition 8. Let $F, G \in H \otimes H$ be twists on a QBA H . Then the twisted structures induced by F and G coincide if and only if there exists a compatible twist $C \in H \otimes H$ such that $G = FC$.

Proof. We have already seen that if F, G give rise to the same QBA structure then $C = F^{-1}G$ is a compatible twist and $G = FC$. Conversely, suppose C is a compatible twist and set $G = FC$. Then

$$\begin{aligned}\Delta_G &= \Delta_{FC} = (\Delta_C)_F = \Delta_F \\ \Phi_G &= \Phi_{FC} = (\Phi_C)_F = \Phi_F\end{aligned}$$

so that G gives precisely the same twisted structure as F . □

Setting $G = 1 \otimes 1$ into the above gives

Corollary. Let $F \in H \otimes H$ be a twist on a QBA H . Then the twisted structure induced by F coincides with the structure on H if and only if F is a compatible twist.

In view of the group properties of twists the above corollary is equivalent to proposition 8.

Let H be a quasi-triangular QHA with R -matrix $\mathcal{R}$ satisfying equation (6.31). From proposition 7, the opposite co-associator $\Phi^T = \Phi_{321}^{-1}$ and co-product Δ^T are obtained by twisting with $\mathcal{R}$, so that $\Phi^T = \Phi_{\mathcal{R}}$. The proof of this result utilises only the properties (6.31). Hence, since

$$\Phi = \Phi_{\mathcal{R}^{-1}\mathcal{R}} = (\Phi_{\mathcal{R}})_{\mathcal{R}^{-1}} = (\Phi^T)_{\mathcal{R}^{-1}}$$

it follows that if Q is another R -matrix for H i.e. satisfies equation (6.31), then we must have also

$$(\Phi^T)_{Q^{-1}} = \Phi.$$

Then $Q^{-1}\mathcal{R}$ must qualify as a compatible twist. Indeed it obviously commutes with Δ , while as to the quasi-cocycle condition, we have

$$\Phi_{Q^{-1}\mathcal{R}} = (\Phi_{\mathcal{R}})_{Q^{-1}} = (\Phi^T)_{Q^{-1}} = \Phi.$$

Note that $(Q^T)^{-1}, (\mathcal{R}^T)^{-1}$ also determine R -matrices so the following must all determine compatible twists: $Q^{-1}\mathcal{R}, Q^T\mathcal{R}, \mathcal{R}^{-1}Q, \mathcal{R}^TQ$. In particular $\mathcal{R}^T\mathcal{R}$ must determine a compatible twist, as may be verified directly.

With the notation of section 4, it is easily seen that the operator

$$A = \Delta(u^{-1})F_{\delta}^{-1}(u \otimes u)F_0 = F_{\delta}^{-1}(u \otimes u)F_0\Delta(u^{-1}) \quad (7.42)$$

commutes with Δ . This operator appears in the work of Altschuler and Coste [1] in connection with ribbon QHAs. The operator A satisfies the quasi-cocycle condition and thus determines a compatible twist.

For general QBAs H , to see that there are sufficiently many compatible twists, we have

Lemma 4. *Let $z \in H$ be an invertible central element. Then*

$$C = (z \otimes z)\Delta(z^{-1})$$

is a compatible twist.

Proof. Obviously C commutes with the co-product Δ so it remains to prove that it satisfies the quasi-cocycle condition. To this end note that

$$\begin{aligned} (C \otimes 1)(\Delta \otimes 1)C &= (z \otimes z \otimes 1)(\Delta(z^{-1}) \otimes 1)(\Delta(z) \otimes z)(\Delta \otimes 1)\Delta(z^{-1}) \\ &= (z \otimes z \otimes z)(\Delta \otimes 1)\Delta(z^{-1}) \end{aligned} \quad (7.43)$$

and similarly

$$\begin{aligned} (1 \otimes C)(1 \otimes \Delta)C &= (1 \otimes z \otimes z)(1 \otimes \Delta(z^{-1}))(z \otimes \Delta(z))(1 \otimes \Delta)\Delta(z^{-1}) \\ &= (z \otimes z \otimes z)(1 \otimes \Delta)\Delta(z^{-1}) \end{aligned} \quad (7.44)$$

thus

$$\begin{aligned} (C \otimes 1)(\Delta \otimes 1)C\Phi &\stackrel{(7.43)}{=} (z \otimes z \otimes z)(\Delta \otimes 1)\Delta(z^{-1})\Phi \\ &\stackrel{(2.1)}{=} (z \otimes z \otimes z)\Phi(1 \otimes \Delta)\Delta(z^{-1}) \\ &\stackrel{(7.44)}{=} (z \otimes z \otimes z)\Phi(z^{-1} \otimes z^{-1} \otimes z^{-1}) \\ &\quad (1 \otimes C)(1 \otimes \Delta)C \\ &= \Phi(1 \otimes C)(1 \otimes \Delta)C. \end{aligned}$$

□

With C as in the lemma, we see that

$$(\epsilon \otimes 1)C = (1 \otimes \epsilon)C = \epsilon(z).$$

Thus, strictly speaking, $\epsilon(z^{-1})C$ qualifies as a compatible twist.

Following Altschuler and Coste [1], a quasi-triangular QHA is called a ribbon QHA if the operator A of equation (7.42) is given by

$$A = (v \otimes v)\Delta(v^{-1})$$

for a certain invertible central element v , related to the u -operator u . This is consistent with the lemma above and the fact that A determines a compatible twist.

In the case of ribbon Hopf algebras, we have $\mathcal{R}^T \mathcal{R} = (v \otimes v)\Delta(v^{-1})$, so that the compatible twist $\mathcal{R}^T \mathcal{R}$ is also of the form of lemma 4. This may not be the case for quasi-triangular QHAs in general.

It is worth noting that if H is a QHA and $C \in H \otimes H$ a compatible twist then H is also a QHA under the twisted structure induced by C with exactly the same co-product Δ , co-unit ϵ , co-associator Φ , antipode S , but with canonical elements given by equation (4.17); viz

$$\alpha_C = m \cdot (S \otimes 1)(1 \otimes \alpha)C^{-1}, \quad \beta_C = m \cdot (1 \otimes S)(1 \otimes \beta)C.$$

In view of Theorem 1 and its corollary, we have immediately

Proposition 9. *Suppose $C \in H \otimes H$ is a compatible twist on a QHA H . Then there exists a unique invertible central element $z \in H$ such that*

$$z\alpha = \alpha_C, \quad \beta_C z = \beta.$$

Explicitly

$$\begin{aligned} z &= S(X_\nu)\alpha_C Y_\nu \beta S(Z_\nu) = \bar{X}_\nu \beta S(\bar{Y}_\nu)\alpha_C \bar{Z}_\nu \\ z^{-1} &= S(X_\nu)\alpha Y_\nu \beta_C S(Z_\nu) = \bar{X}_\nu \beta_C S(\bar{Y}_\nu)\alpha \bar{Z}_\nu. \end{aligned}$$

In the case H is quasi-triangular we have seen that $C = \mathcal{R}^T \mathcal{R}$ is a compatible twist. Since the latter form a group we have the infinite family of compatible twists $C = (\mathcal{R}^T \mathcal{R})^m$, $m \in \mathbb{Z}$, in which case the central elements $z^{\pm 1}$ of proposition 9 give the quadratic invariants of [12].

We conclude this section by noting, in the quasi-triangular case, that twisting the Drinfeld twist with the R -matrix $\mathcal{R}$ gives, from Theorem 4, the twisted Drinfeld twist

$$F_\delta^{\mathcal{R}} \equiv (F_\delta)_{\mathcal{R}} = (S \otimes S)(\mathcal{R}^T)^{-1} \cdot F_\delta \cdot \mathcal{R}^{-1}.$$

On the other hand, since $(\mathcal{R}^T)^{-1}$ is an R -matrix we have, from eq. (6.33),

$$(S \otimes S)(\mathcal{R}^T)^{-1} = F_\delta^T (\mathcal{R}^T)^{-1} F_\delta^{-1}$$

which implies

$$F_\delta^{\mathcal{R}} = F_\delta^T (\mathcal{R}^T)^{-1} \cdot \mathcal{R}^{-1} = F_\delta^T (\mathcal{R} \mathcal{R}^T)^{-1}$$

where $\mathcal{R} \mathcal{R}^T$ and its inverse are compatible twists under the opposite structure. This shows that F_δ^T will give rise to a Drinfeld twist under the opposite structure of proposition 7 induced by twisting with $\mathcal{R}$ (which has antipode S rather than S^{-1}). Applying T to the equation above gives

$$(F_\delta^{\mathcal{R}})^T = F_\delta (\mathcal{R}^T \mathcal{R})^{-1}$$

which shows that, since $\mathcal{R}^T \mathcal{R}$ and its inverse are compatible twists, $(F_\delta^{\mathcal{R}})^T$ also gives rise to a Drinfeld twist on H .

8 Quasi-dynamical QYBE

Throughout we assume H is a quasi-triangular QHA with R -matrix $\mathcal{R}$ satisfying (6.31) which we reproduce here:

$$\begin{aligned} (i) \quad & \Delta^T(a)\mathcal{R} = \mathcal{R}\Delta(a), \quad \forall a \in H \\ (ii) \quad & (\Delta \otimes 1)\mathcal{R} = \Phi_{231}^{-1} \mathcal{R}_{13} \Phi_{132} \mathcal{R}_{23} \Phi_{123}^{-1} \\ (iii) \quad & (1 \otimes \Delta)\mathcal{R} = \Phi_{312} \mathcal{R}_{13} \Phi_{213}^{-1} \mathcal{R}_{12} \Phi_{123}. \end{aligned} \tag{6.31'}$$

Applying $T \otimes 1$ to (ii) and $1 \otimes T$ to (iii) then gives

$$\begin{aligned} (ii') \quad & (\Delta^T \otimes 1)\mathcal{R} = \Phi_{321}^{-1} \mathcal{R}_{23} \Phi_{312} \mathcal{R}_{13} \Phi_{213}^{-1} \\ (iii') \quad & (1 \otimes \Delta^T)\mathcal{R} = \Phi_{321} \mathcal{R}_{12} \Phi_{231}^{-1} \mathcal{R}_{13} \Phi_{132}. \end{aligned}$$

It follows that

$$\mathcal{R}_{12}(\Delta \otimes 1)\mathcal{R} = (\Delta^T \otimes 1)\mathcal{R} \cdot \mathcal{R}_{12}$$

from which we deduce that $\mathcal{R}$ must satisfy the quasi-QYBE:

$$\mathcal{R}_{12}\Phi_{231}^{-1}\mathcal{R}_{13}\Phi_{132}\mathcal{R}_{23}\Phi_{123}^{-1} = \Phi_{321}^{-1}\mathcal{R}_{23}\Phi_{312}\mathcal{R}_{13}\Phi_{213}^{-1}\mathcal{R}_{12}. \quad (8.45)$$

If we twist H with a twist $F \in H \otimes H$ then H is also a quasi-triangular QHA under the twisted structure (4.16,4.17) induced by F with universal R -matrix

$$\mathcal{R}_F = F^T \mathcal{R} F^{-1}.$$

Following equation (7.40) we say a twist $F(\lambda) \in H \otimes H$ satisfies the shifted quasi-cocycle condition if

$$[F(\lambda) \otimes 1] \cdot (\Delta \otimes 1)F(\lambda) \cdot \Phi = \Phi \cdot [1 \otimes F(\lambda + h^{(1)})] \cdot (1 \otimes \Delta)F(\lambda) \quad (8.46)$$

where $\lambda \in H$ depends on one (or possibly several) parameters and $h \in H$ is fixed. Alternatively, we may write in obvious notation

$$F_{12}(\lambda) \cdot (\Delta \otimes 1)F(\lambda) \cdot \Phi = \Phi \cdot F_{23}(\lambda + h^{(1)}) \cdot (1 \otimes \Delta)F(\lambda). \quad (8.46')$$

When $h = 0$, this reduces to the quasi-cocycle condition (7.40) satisfied by $F = F(\lambda)$. When $\Phi = 1 \otimes 1 \otimes 1$ (i.e. the normal Hopf-algebra case) equation (8.46) reduces to the usual shifted cocycle condition.

Twisting H with a twist F satisfying the (unshifted) quasi-cocycle condition results in a QHA with the same co-associator Φ , co-unit ϵ and antipode S but with the twisted co-product Δ_F , R -matrix $\mathcal{R}_F$ (and canonical elements α_F, β_F). We now consider twisting H with a twist $F = F(\lambda)$ satisfying the shifted condition (8.46). Then under this twisted structure H is also a quasi-triangular QHA with the same co-unit ϵ and antipode S but with the co-associator $\Phi(\lambda) = \Phi_{F(\lambda)}$, and co-product and R -matrix given by

$$\Delta_\lambda(a) = F(\lambda)\Delta(a)F(\lambda)^{-1}, \quad \forall a \in H, \quad \mathcal{R}(\lambda) = F^T(\lambda)\mathcal{R}F(\lambda)^{-1} \quad (8.47)$$

with canonical elements $\alpha_\lambda = \alpha_{F(\lambda)}, \beta_\lambda = \beta_{F(\lambda)}$.

In view of equation (8.46') we have for the co-associator

$$\begin{aligned} \Phi(\lambda) &= F_{12}(\lambda) \cdot (\Delta \otimes 1)F(\lambda) \cdot \Phi \cdot (1 \otimes \Delta)F(\lambda)^{-1} \cdot F_{23}(\lambda)^{-1} \\ &= \Phi \cdot F_{23}(\lambda + h^{(1)}) \cdot (1 \otimes \Delta)F(\lambda) \cdot (1 \otimes \Delta)F(\lambda)^{-1} \cdot F_{23}(\lambda)^{-1} \\ &= \Phi \cdot F_{23}(\lambda + h^{(1)}) \cdot F_{23}(\lambda)^{-1} \end{aligned} \quad (8.48)$$

which implies

$$\Phi(\lambda)^{-1} = F_{23}(\lambda) \cdot F_{23}(\lambda + h^{(1)})^{-1} \cdot \Phi^{-1}.$$

In the Hopf-algebra case equation (8.48) reduces to the expression for $\Phi(\lambda)$ obtained in [13] ($\Phi = 1 \otimes 1 \otimes 1$).

Under the above twisted structure equation (6.31)(ii) becomes

$$(\Delta_\lambda \otimes 1)\mathcal{R}(\lambda) = \Phi_{231}(\lambda)^{-1} \cdot \mathcal{R}_{13}(\lambda) \cdot \Phi_{132}(\lambda) \cdot \mathcal{R}_{23}(\lambda) \cdot \Phi_{123}^{-1}(\lambda).$$

Now

$$\begin{aligned} \Phi_{132}(\lambda) &= (1 \otimes T)\Phi_{123}(\lambda) \\ &\stackrel{(8.48)}{=} \Phi_{132} \cdot F_{23}^T(\lambda + h^{(1)}) \cdot F_{23}^T(\lambda)^{-1} \end{aligned} \quad (8.49)$$

which implies

$$\begin{aligned}
(\Delta_\lambda \otimes 1)\mathcal{R}(\lambda) &= \Phi_{231}(\lambda)^{-1} \cdot \mathcal{R}_{13}(\lambda) \cdot \Phi_{132} \cdot F_{23}^T(\lambda + h^{(1)}) \\
&\quad \cdot F_{23}^T(\lambda)^{-1} \cdot \mathcal{R}_{23}(\lambda) \cdot \Phi_{123}^{-1}(\lambda) \\
&\stackrel{(8.48)}{=} \Phi_{231}(\lambda)^{-1} \cdot \mathcal{R}_{13}(\lambda) \cdot \Phi_{132} \cdot F_{23}^T(\lambda + h^{(1)}) \\
&\quad \cdot F_{23}^T(\lambda)^{-1} \cdot \mathcal{R}_{23}(\lambda) \cdot F_{23}(\lambda) \cdot F_{23}(\lambda + h^{(1)})^{-1} \cdot \Phi_{123}^{-1} \\
&\stackrel{(8.47)}{=} \Phi_{231}(\lambda)^{-1} \cdot \mathcal{R}_{13}(\lambda) \cdot \Phi_{132} \cdot \mathcal{R}_{23}(\lambda + h^{(1)}) \cdot \Phi_{123}^{-1}.
\end{aligned}$$

Similarly equation (6.31)(iii) becomes

$$(1 \otimes \Delta_\lambda)\mathcal{R}(\lambda) = \Phi_{312}(\lambda) \cdot \mathcal{R}_{13}(\lambda) \cdot \Phi_{213}^{-1}(\lambda) \cdot \mathcal{R}_{12}(\lambda) \cdot \Phi_{123}(\lambda).$$

Now

$$\begin{aligned}
\Phi_{312}(\lambda) &= (T \otimes 1)(1 \otimes T)\Phi_{123}(\lambda) \\
&\stackrel{(8.48)}{=} (T \otimes 1)[\Phi_{132} \cdot F_{23}^T(\lambda + h^{(1)}) \cdot F_{23}^T(\lambda)^{-1}] \\
&= \Phi_{312} \cdot F_{13}^T(\lambda + h^{(2)}) \cdot F_{13}^T(\lambda)^{-1}
\end{aligned}$$

while

$$\begin{aligned}
\Phi_{213}^{-1}(\lambda) &= (T \otimes 1)\Phi(\lambda)^{-1} \\
&\stackrel{(8.48)}{=} (T \otimes 1)[F_{23}(\lambda) \cdot F_{23}(\lambda + h^{(1)})^{-1} \cdot \Phi^{-1}] \\
&= F_{13}(\lambda) \cdot F_{13}(\lambda + h^{(2)})^{-1} \cdot \Phi_{213}^{-1}.
\end{aligned}$$

Therefore

$$\begin{aligned}
(1 \otimes \Delta_\lambda)\mathcal{R}(\lambda) &= \Phi_{312} \cdot F_{13}^T(\lambda + h^{(2)}) \cdot F_{13}^T(\lambda)^{-1} \cdot \mathcal{R}_{13}(\lambda) \\
&\quad \cdot F_{13}(\lambda) \cdot F_{13}(\lambda + h^{(2)})^{-1} \cdot \Phi_{213}^{-1} \cdot \mathcal{R}_{12}(\lambda) \cdot \Phi_{123}(\lambda) \\
&\stackrel{(8.47)}{=} \Phi_{312} \cdot \mathcal{R}_{13}(\lambda + h^{(2)}) \cdot \Phi_{213}^{-1} \cdot \mathcal{R}_{12}(\lambda) \cdot \Phi_{123}(\lambda).
\end{aligned}$$

We thus arrive at

Lemma 5. $\mathcal{R}(\lambda)$ satisfies the co-product properties

$$\begin{aligned}
(i) \quad & (\Delta_\lambda \otimes 1)\mathcal{R}(\lambda) = \Phi_{231}^{-1}(\lambda) \cdot \mathcal{R}_{13}(\lambda) \cdot \Phi_{132} \cdot \mathcal{R}_{23}(\lambda + h^{(1)}) \cdot \Phi_{123}^{-1} \\
(ii) \quad & (1 \otimes \Delta_\lambda)\mathcal{R}(\lambda) = \Phi_{312} \cdot \mathcal{R}_{13}(\lambda + h^{(2)}) \cdot \Phi_{213}^{-1} \cdot \mathcal{R}_{12}(\lambda) \cdot \Phi_{123}(\lambda) \\
(iii) \quad & (\Delta_\lambda^T \otimes 1)\mathcal{R}(\lambda) = \Phi_{321}^{-1}(\lambda) \cdot \mathcal{R}_{23}(\lambda) \cdot \Phi_{312} \cdot \mathcal{R}_{13}(\lambda + h^{(2)}) \cdot \Phi_{213}^{-1} \\
(iv) \quad & (1 \otimes \Delta_\lambda^T)\mathcal{R}(\lambda) = \Phi_{321} \cdot \mathcal{R}_{12}(\lambda + h^{(3)}) \cdot \Phi_{231}^{-1} \cdot \mathcal{R}_{13}(\lambda) \cdot \Phi_{132}(\lambda).
\end{aligned} \tag{8.50}$$

Proof. We have already proved (i) and (ii) while (iii) follows by applying $(T \otimes 1)$ to (i) and (iv) by applying $(1 \otimes T)$ to (ii). $\square$

We are now in a position to determine the QQYBE (8.45) satisfied by $\mathcal{R} = \mathcal{R}(\lambda)$ for this twisted structure. We have

$$\begin{aligned}
& \mathcal{R}_{23}(\lambda) \cdot \Phi_{312} \cdot \mathcal{R}_{13}(\lambda + h^{(2)}) \cdot \Phi_{213}^{-1} \cdot R_{12}(\lambda) \\
\stackrel{(8.50)(ii)}{=} & \mathcal{R}_{23}(\lambda) \cdot (1 \otimes \Delta_\lambda) \mathcal{R}(\lambda) \cdot \Phi_{123}^{-1}(\lambda) \\
\stackrel{(6.31)(i)}{=} & (1 \otimes \Delta_\lambda^T) \mathcal{R}(\lambda) \cdot \mathcal{R}_{23}(\lambda) \cdot \Phi_{123}^{-1}(\lambda) \\
\stackrel{(8.50)(iv)}{=} & \Phi_{321} \cdot R_{12}(\lambda + h^{(3)}) \cdot \Phi_{231}^{-1} \cdot R_{13}(\lambda) \cdot \Phi_{132}(\lambda) \cdot \mathcal{R}_{23}(\lambda) \cdot \Phi_{123}^{-1}(\lambda)
\end{aligned}$$

where for the last three terms we have

$$\begin{aligned}
\Phi_{132}(\lambda) \cdot \mathcal{R}_{23}(\lambda) \cdot \Phi_{123}^{-1}(\lambda) & \stackrel{(8.48, 8.49)}{=} \Phi_{132} \cdot F_{23}^T(\lambda + h^{(1)}) \cdot F_{23}^T(\lambda)^{-1} \\
& \cdot R_{23}(\lambda) \cdot F_{23}(\lambda) \cdot F_{23}(\lambda + h^{(1)})^{-1} \cdot \Phi_{123}^{-1} \\
& \stackrel{(8.47)}{=} \Phi_{132} \cdot \mathcal{R}_{23}(\lambda + h^{(1)}) \cdot \Phi_{123}^{-1}.
\end{aligned}$$

Hence

$$\begin{aligned}
& \mathcal{R}_{23}(\lambda) \cdot \Phi_{312} \cdot \mathcal{R}_{13}(\lambda + h^{(2)}) \cdot \Phi_{213}^{-1} \cdot R_{12}(\lambda) \\
= & \Phi_{321} \cdot \mathcal{R}_{12}(\lambda + h^{(3)}) \cdot \Phi_{231}^{-1} \cdot \mathcal{R}_{13}(\lambda) \cdot \Phi_{132} \cdot \mathcal{R}_{23}(\lambda + h^{(1)}) \cdot \Phi_{123}^{-1}.
\end{aligned}$$

We thus arrive at

Proposition 10. $\mathcal{R}(\lambda)$ satisfies the quasi-dynamical QYBE

$$\begin{aligned}
& \mathcal{R}_{12}(\lambda + h^{(3)}) \cdot \Phi_{231}^{-1} \cdot \mathcal{R}_{13}(\lambda) \cdot \Phi_{132} \cdot \mathcal{R}_{23}(\lambda + h^{(1)}) \cdot \Phi_{123}^{-1} \\
= & \Phi_{321}^{-1} \cdot \mathcal{R}_{23}(\lambda) \cdot \Phi_{312} \cdot \mathcal{R}_{13}(\lambda + h^{(2)}) \cdot \Phi_{213}^{-1} \cdot \mathcal{R}_{12}(\lambda).
\end{aligned} \tag{8.51}$$

In the Hopf algebra case ($\Phi = 1 \otimes 1 \otimes 1$) equation (8.51) reduces to the usual dynamical QYBE. If we set $h = 0$ then equation (8.51) reduces to the quasi-QYBE (8.45) satisfied by $\mathcal{R} = \mathcal{R}(\lambda)$. Hence the term quasi-dynamical QYBE for (8.51): we could, alternatively, refer to (8.51) as the dynamical quasi-QYBE (dynamical QQYBE), since it is obviously the quasi-Hopf algebra analogue of the usual dynamical QYBE.

With respect to the QHA structure of propositions 2, 2' we have the R -matrices

$$\mathcal{R}'(\lambda) = (S \otimes S) \mathcal{R}(\lambda), \quad \mathcal{R}_0(\lambda) = (S^{-1} \otimes S^{-1}) \mathcal{R}(\lambda)$$

respectively. Then applying $(S \otimes S \otimes S)$, $(S^{-1} \otimes S^{-1} \otimes S^{-1})$ respectively to equation (8.51) it follows that both of these R -matrices satisfy the opposite quasi-dynamical QYBE

$$\begin{aligned}
& \tilde{\mathcal{R}}_{12}(\lambda) \cdot \tilde{\Phi}_{231}^{-1} \cdot \tilde{\mathcal{R}}_{13}(\lambda + h^{(2)}) \cdot \tilde{\Phi}_{132} \cdot \tilde{\mathcal{R}}_{23}(\lambda) \cdot \tilde{\Phi}_{123}^{-1} \\
= & \tilde{\Phi}_{321}^{-1} \cdot \tilde{\mathcal{R}}_{23}(\lambda + h^{(1)}) \cdot \tilde{\Phi}_{312} \cdot \tilde{\mathcal{R}}_{13}(\lambda) \cdot \tilde{\Phi}_{213}^{-1} \cdot \tilde{\mathcal{R}}_{12}(\lambda + h^{(3)})
\end{aligned}$$

where $\tilde{\Phi}$ is the co-associator of propositions 2, 2' and $\tilde{\mathcal{R}}(\lambda)$ denotes $\mathcal{R}'(\lambda)$, $\mathcal{R}_0(\lambda)$ respectively. Moreover applying $(T \otimes 1)((1 \otimes T)(T \otimes 1))$ to equation (8.51) it is easily seen that $\mathcal{R}^T(\lambda)$ also satisfies the above opposite quasi-dynamical QYBE but with respect to the opposite co-associator Φ^T of proposition 1.

We anticipate that the quasi-dynamical QYBE will play an important role in obtaining elliptic solutions to the QQYBE from trigonometric ones via twisted QYEs. Of particular interest is the quasi-dynamical QYBE for elliptic quantum groups.

References

- [1] Altschuler, D., Coste, A.: *Quasi-quantum groups, knots, three manifolds and topological field theory*, Commun. Math. Phys., **150** No. 1 83–107 (1992) [arXiv:hep-th/9202047](#).
- [2] Chari, V., Pressley, A.: *A Guide to Quantum Groups*, Cambridge University Press, Cambridge (1994).
- [3] Dijkgraaf, R., Pasquier, V., Roche, P.: *Quasi-quantum groups related to orbifold models*, Proc. Int. Coll. on Modern Quantum Field Theory, Tata Institute, Bombay, (1990).
- [4] Dijkgraaf, R., Vafa, C., Verlinde, E., Verlinde, H.: *The operator algebra of orbifold models*, Commun. Math. Phys., **123** No. 3 485–526 (1989).
- [5] Drinfeld, V.G.: *On almost cocommutative Hopf algebras*, Leningrad Math. J., **1** No. 2 321–342 (1990).
- [6] Drinfeld, V.G.: *Quasi-Hopf Algebras*, Leningrad Math. J., **1** No. 6 1419–1457 (1990).
- [7] Drinfeld, V.G.: *On quasitriangular quasi-Hopf algebras and a group closely connected with $Gal(\bar{\mathbb{Q}}/\mathbb{Q})$* , Leningrad Math. J., **2** No. 4 829–860 (1991).
- [8] Enriquez, B., Felder, G.: *Elliptic quantum group $E_{\tau, \eta}(\mathfrak{sl}_2)$ and quasi-Hopf algebras*, Commun. Math. Phys., **195** 651–689, (1998) [arXiv:q-alg/9703018](#).
- [9] Foda, O., Iohara, K., Jimbo, M., Kedem, R., Miwa, T., Yan, H.: *An elliptic quantum algebra for $\widehat{\mathfrak{sl}}_2$* , Lett. Math. Phys., **32** No. 3 259–268, (1994) [arXiv:hep-th/9403094](#).
- [10] Felder, G.: *Elliptic quantum groups*, Proc. XIth ICMP (Paris, 1994), 211–218, (1995).
- [11] Fronsdal, C.: *Quasi-Hopf deformations of quantum groups*, Lett. Math. Phys., **40** 117–134, (1997) [arXiv:q-alg/9611028](#).
- [12] Gould, M.D., Zhang, Y.-Z., Isaac, P.S.: *Casimir invariants from quasi-Hopf (super) algebras*, J. Math. Phys., **41** 547–568, (2000) [arXiv:math.QA/9811052](#).
- [13] Gould, M.D., Zhang, Y.-Z., Isaac, P.S.: *On quasi-Hopf superalgebras*, Commun. Math. Phys., **224** 341–372 (2001) [arXiv:math.QA/9811062](#).
- [14] Jimbo M., Odake S., Konno H., Shiraishi J.: *Quasi-Hopf twistors for elliptic quantum groups*, Transform. Groups, **4** 303–327, (1999) [arXiv:q-alg/9712029](#).
- [15] Links, J. R., Gould, M. D., Zhang, Y.-Z.: *Twisting invariance of link polynomials derived from ribbon quasi-Hopf algebras*, J. Math. Phys., **41**, 5020–5032, (2000) [arXiv:math.QA/9904069](#).
- [16] Mack, G., Schomerus, V.: *Quasi quantum group symmetry and local braid relations in the conformal Ising model*, Phys. Lett. B, **267**, 207–213, (1991).
- [17] Majid, S.: *Foundations of Quantum Group Theory*, Cambridge University Press, Cambridge (2000).
- [18] Reshetikhin, N.: *Quasitriangle Hopf algebras and invariants of links*, Leningrad Math. J., **1** No. 2 491–513 (1990).
- [19] Sweedler, M.E.: *Hopf Algebras*, Benjamin, New York (1969).

- [20] Zhang, Y.-Z., Gould, M.D.: *Quasi-Hopf superalgebras and elliptic quantum supergroups*, J. Math. Phys., **40** No. 10 5264–5282 (1999) [arXiv:math.QA/9809156](#).